



# Coronavirus Disease 2019 (COVID-19)

## National Surveillance Report as of 12/05/2020

Date of publication: 14/05/2020

Produced by the Epidemiological Surveillance Unit of the Ministry of Health

Contributors: Annalisa Quattrocchi, Ioannis Mamais, Valentinos Silvestros, Anna Demetriou (Health Monitoring Unit), Maria Athanasiadou (Health Monitoring Unit), Theopisti Kyprianou (Health Monitoring Unit), Androulla Stylianou, Sotiroula Sotiriou, Fani Theofhanous, Christos Charalambous, Ioanna Gregoriou, Maria Koliou, Georgios Nikolopoulos and Elisavet Constantinou

Scientific Committee: Elisavet Constantinou, Constantinos Constantinou, Niki Paphitou, Georgios Nikolopoulos, Maria Koliou, George Panos, Eirini Christaki, Zoi - Dorothea Pana, Constantinos Tsioutis, Leondios Kostrikis, Peter Karayiannis, George Petrikkos, Petros Agathangelou, George Mixides

Suggested citation: Epidemiological Surveillance Unit of the Ministry of Health, Cyprus. National Situation Report. Coronavirus Disease 2019 (COVID-19), 14 May 2020.



## Summary

- As of May 12<sup>th</sup>, a total of 904 COVID-19 cases and 23 deaths (case fatality rate: 2.5%) have been reported in the Republic of Cyprus.
- Among these cases, 20.9% are health-care workers (n = 189) - 4.3% physicians (n = 39), 10.5% nurses (n = 95), 1.4% other health occupations (n = 13), and 4.7% auxiliary staff (n = 42).
- The median age of cases is 45 years (interquartile range - IQR: 32-59 years); 49.8% are female and 50.2% are male.
- Overall, of 760 cases for which the place of exposure was known, locally acquired infections (index cases and close-contacts of confirmed cases) were 631 (83%) - of these 8.2% (n = 62) were related to a health-care facility (General Hospital in Pafos) and 12.7% (n = 80) were reported in Aradippou municipality.
- In total, 19.1% (n = 173) of cases received hospital care, of which 135 (78%) have been discharged alive from the hospital. Median age of all hospitalized patients is: 62 years (interquartile range: 49-73 years), and 65.3% are males.
- Four patients were still in intensive care units (for part of the day if they died, were discharged or transferred on that day or for the whole day, by May 12<sup>th</sup>); all of these were intubated.
- Overall, 481 (53.2%) cases have recovered.
- A total of 78,239 tests have been performed as of May 12<sup>th</sup> (8,932.4 per 100,000 population).



## Epidemiological surveillance in the Republic of Cyprus

Analyses are based on laboratory-confirmed cases notified to the Epidemiological Surveillance Unit of the Ministry of Health.

As of May 12<sup>th</sup>, 904 laboratory-confirmed cases of coronavirus disease 2019 (COVID-19) have been reported (Figure 1 and 2).

The median time between symptoms onset and date of sampling was 4 days (Interquartile range - IQR: 2-7 days). It should be noted that for 14 cases the date of sample collection was before the onset of symptoms because of immediate testing of contacts of possible and laboratory-confirmed cases.

As of May 12<sup>th</sup>, the 14-day cumulative incidence rate of COVID-19 (per 100,000 population), a measure which reflects the number of active COVID-19 cases in the population (prevalence)<sup>1</sup>, is 7 per 100,000 population (Figure 3).




Figure 1: Number of laboratory-confirmed COVID-19-cases in Cyprus since 01/03/2020 by date of sample collection and date of symptoms onset (n = 904 and n = 477 with data available, respectively).

*Recent data should be interpreted with caution due to the possibility that cases with date of onset within the reporting period have not yet been diagnosed.*

<sup>1</sup>Coronavirus disease 2019 (COVID-19) pandemic: increased transmission in the EU/EEA and the UK – seventh update, 25 March 2020. Stockholm: ECDC; 2020.

<https://www.ecdc.europa.eu/sites/default/files/documents/RRA-seventh-update-Outbreak-of-coronavirus-disease-COVID-19.pdf>




Figure 2: Number and cumulative number of laboratory-confirmed COVID-19-cases in Cyprus since 01/03/2020, by date of laboratory reporting (n = 904). Recent data should be interpreted with caution due to the possibility that cases with date of onset within the reporting period have not yet been diagnosed.




Figure 3. COVID-19 14-day cumulative incidence rate per 100,000 population (proxy of COVID-19 prevalence). March 23<sup>rd</sup> represents the first 14<sup>th</sup> day since cases have been reported.

## Characteristics of the cases

Among these cases, 50.2% are male (n = 454) and 49.8% female (n = 450).

The median age of cases is 45 years (IQR: 32-59 years). By age groups, cases included 56 infants, children and adolescents aged 0-17 years-old (6.2%), 625 adults aged 18-59 years (69.1%), and 223 persons aged 60 years and older (24.7%) (Figure 4).




Figure 4: Laboratory-confirmed COVID-19-cases in Cyprus by sex and age groups.

Among all cases, 350 (38.7%) were reported in Nicosia district, 234 (25.9%) in Larnaka, 155 (17.2%) in Pafos, 103 (11.4%) in Limassol, 41 (4.5%) in Ammochostos, and 21 (2.3%) were reported either in British bases or had a residence abroad, or information was not available (Table A1 in appendix).

Figure A1 in appendix shows the distribution of cases by postal code.

Notably, 123 cases (13.6%) were reported in Aradippou, a municipality in Larnaka district (Table A1 in appendix). Cases in Aradippou, including a cluster in a local bakery production line, are mainly males (58.5%; n = 72) and the median age is 49 years (IQR: 33-62 years). If the cluster is excluded, cases are mainly female (53.7%; n = 51) and the median age is 55 years (IQR: 39-69years).



Among the 904 cases, 20.9% are health-care workers<sup>2</sup> (n = 189) - 4.3% physicians (n = 39), 10.5% nurses (n = 95), 1.4% other health occupations (n = 13), and 4.7% auxiliary staff (n = 42).

Table 1 shows the distribution of health-care workers by district of residence.

Table 1: Health-care workers by district of residence (n=189).

District	Health-care worker	Physicians	Nurses	Other health occupations	Auxiliary staff
Ammochostos	16	3	7	1	5
Larnaka	43	7	24	3	9
Limassol	16	3	9	2	2
Nicosia	55	12	23	5	15
Pafos	59	14	32	2	11
Total	189	39	95	13	42

### Epidemiological link

As of May 12<sup>th</sup>, place of exposure is available for 760 cases (84.1%).

In total, 17% (n = 129) of laboratory-confirmed COVID-19-cases had history of travel or residence abroad during the 14 days prior to symptom onset (imported). These cases have a direct link to the UK and Greece, mainly.

Locally acquired infections (index cases and close-contacts of confirmed cases) occurred in 83% (n = 631 of 760 with known place of exposure) of the cases, of which 8.2% (n = 62) were related to a health-care facility (General Hospital in Pafos).

Of all cases in Aradippou (Larnaka district) (n = 123), 80 (65%) were locally-acquired, 10 (8.1%) imported and for 33 cases (26.8%) the epidemiological link was not recorded at the moment.

Table A1 in the appendix shows the number and the rate (per 100,000 population) of locally-acquired cases by district of residence.

<sup>2</sup> The term "health-care worker" is based on the occupation and not on the place of exposure. Health-care workers are defined as all health care professionals, allied health workers, and auxiliary health workers.



## Clinical features

Of the 904 laboratory-confirmed COVID-19-cases, clinical information is available for 98.5% (n = 890), of which 31.1% (n = 269) reported no symptoms at diagnosis and 67.5% (n = 610) reported at least one symptom. The most commonly reported symptoms were:

- cough (308/879; 35%),
- fever (284/878; 32.3%),
- myalgia (199/877; 22.7%),
- sore throat (156/875; 17.8%),
- anosmia (107/787; 13.6%), and
- shortness of breath (105/862; 12.2%).

Other reported symptoms were diarrhoea, runny nose, and headache.

Table A2 in appendix reports the sex and age distribution of asymptomatic cases at diagnosis.

## Pre-existing conditions

Information on comorbidities was available for 795 (87.9%) cases. Of these, 325 (40.8%) reported at least one comorbidity.

The most commonly reported comorbidities were:

- hypertension (126/788; 16%),
- diabetes (72/769; 9.4%),
- heart disease (62/790; 7.8%), and
- cancer (24/795; 3%).

Other reported comorbidities were chronic kidney disease, autoimmune disease, and chronic respiratory disease.



## Deaths

As of May 12<sup>th</sup>, 23 deaths were reported in Cyprus (Case Fatality Rate - CFR: 2.5%). The mortality rate for COVID-19 is 2.6 per 100,000 population.

Seventeen deaths (73.9%) occurred in men and six (26.1%) in women; the median age of all deaths was 76 years (IQR: 66-79 years). Nine deaths were reported among residents in Larnaka, six in Pafos, three in Nicosia and Ammochostos, each, and two in Limassol (Appendix Table A3).

The median time from date of sampling to death was 8 days (IQR: 4-18 days). Figure A3 shows the Kaplan-Meier curve of the time from date of sampling to death.

For 16 deaths, COVID-19 was the underlying cause of death (COVID-19 CFR: 1.8%). Figure 5 reports the number of deaths by date.




Figure 5: Number of deaths among COVID-19 cases in Cyprus by date of death (n = 23).



## Hospitalization and intensive care unit (ICU) admissions<sup>3</sup>

In total, 19.1% (n = 173) of people with COVID-19 received hospital care, and 135 patients (78%) have been discharged alive from the hospital. The median age of hospitalized patients was 62 years (IQR: 49-73 years). Hospitalized cases were mainly males (n = 113; 65.3%).

Figure 6 shows the total number of hospital admissions by date.




Figure 6: Number of laboratory-confirmed COVID-19 cases by date of hospital admission (n = 173).

\* Date of hospital admission; for inpatients hospitalised prior to the beginning of the epidemic, it was replaced with date of sampling.

Overall, 32 cases (18.5% of all hospitalized patients) have been admitted to ICU<sup>4</sup>, of which 4 were still in ICU (as of May 12<sup>th</sup>).

A total of 27 ICU patients (84.4% of all ICU patients) have been intubated, of which 4 (100% of all patients currently in ICU) are still intubated.

The overall median length of stay in ICU (for all 32 ICU cases, considering those still in ICU until May 12<sup>th</sup>) was 11 days (IQR: 8-27.5 days). Figure A4 shows the Kaplan-Meier curve of the length of stay in ICU.

<sup>3</sup> Data on hospitalisation and ICU are provisional and should be interpreted with caution because delay in data reporting is likely; for the construction of the curve, people are no longer in an ICU the day next to the date of their discharge, death or transfer.

<sup>4</sup> Intensive care unit (ICU) refers only to the ICU in Limassol General Hospital and to the ICU in Nicosia General Hospital.



For patients who died while in ICU (n = 15), the median length of stay in ICU was 10 days (IQR: 3-20). Figure A5 shows the Kaplan-Meier curve of the length of stay in ICU for the people who died.

For patients transferred/ discharged alive from ICU (n = 13), the median length of stay in ICU was 10 days (IQR: 8-27 days).

The median age of patients ever admitted to ICU was 65.5 years (IQR: 56-75 years). ICU patients are mainly male (n = 23; 71.9%).

The number of cases currently in ICU is 0.5 per 100,000 population. For comparison, Italy and Lombardia reported the highest rates of 6.7 per 100,000 population (n = 4,068) and 13.8 per 100,000 population (n = 1,381) on April 3rd. The ICU rates in Italy and Lombardia on May 12<sup>th</sup> are 1.6 per 100,000 population (n = 952) and 3.2 per 100,000 population (n = 322)

(<https://github.com/pcm-dpc/COVID-19/blob/master/dati-andamento-nazionale/dpc-covid19-ita-andamento-nazionale-20200512.csv>;

<https://github.com/pcm-dpc/COVID-19/blob/master/dati-regioni/dpc-covid19-ita-regioni-20200512.csv>).

Figure 7 shows the number of patients in ICU, by day and intubation. Table A4 in the appendix shows the total number of ICU admissions by date.




Figure 7: Number of laboratory-confirmed COVID-19 cases in ICU by date and intubation.

*\*Date of discharge/transfer/death included*



## Recovered

As of May 12<sup>th</sup>, 53.2% (n = 481) of COVID-19 cases have recovered<sup>5</sup>.

The median time between the second negative result and the first date of sampling was 21 days (IQR: 18-28 days).

Table 2 shows the number and percentage of recovered cases and their characteristics.

Table 2: Characteristics of recovered cases (n = 481).

Characteristics	Total	Recovered	
	N	n	%
Total	904	481	53.3
Sex			
Male	454	227	50.0
Female	450	254	56.4
Age groups (years)			
0-9	29	11	37.9
10-19	37	18	48.6
20-29	117	64	54.7
30-39	179	105	58.7
40-49	157	77	49.0
50-59	162	92	56.8
60-69	112	59	52.7
70-79	84	43	51.2
80+	27	12	44.4
Median age in years (IQR*)	45 (32-59)	45 (32-59)	

<sup>5</sup> For symptomatic cases, or for cases hospitalised, a COVID-19 case can be considered cured after the resolution of symptoms and two negative tests for SARS-CoV-2 at 24-hour interval at least.

For asymptomatic cases, or for persons isolated at home, the negative tests to document virus clearance should be obtained at a minimum of 14 days after the initial positive test (end of the quarantine period). Novel coronavirus (SARS-CoV-2). Discharge criteria for confirmed COVID-19 cases- When is it safe to discharge COVID-19 cases from the hospital or end home isolation? - Technical Report, 10 March 2020. Stockholm: ECDC; 2020.

## Comparison with selected countries

As of May 12<sup>th</sup>, in Cyprus the reporting rate was 103.2 cases per 100,000 population, the mortality rate was 2.6 deaths per 100,000 population and the CFR was 2.5%.

Table 3 shows COVID-19 indicators for Cyprus and other selected countries.

Figure A2 in appendix reports the rates of cumulative tests and cases (per 100,000 population) in Cyprus and other selected countries. In Cyprus the testing rate is 8,932.4 per 100,000 population.

It should be noted that the number of cases, tests and deaths for Cyprus are aggregated and include people from abroad and the British bases, while the total population does not include inhabitants from abroad or from the British bases.

Table 3: COVID-19 indicators by selected countries, as of 12/05/2020.

Country	N. of cases †	N. of cases (per 100,000 pop)	N. of tests §	N. of tests (per 100,000 pop)	N. of deaths†	CFR <sup>°</sup> (%)	Mortality rate (per 100,000 pop)	Pop. (in thousands)†
Cyprus	904	103.2	78,239	8,932.4	23	2.5	2.6	875.9*
Italy	219,814	363.7	2,673,655	4,424.3	30,739	14.0	50.9	60,431.3
USA	1,347,916	412.0	9,382,235	2,867.7	80,684	6.0	24.7	327,167.4
UK	223,060	335.5	2,007,146	3,018.8	32,065	14.4	48.2	66,488.9
Greece	2,726	25.4	99,363	926.2	151	5.5	1.4	10,727.7
Malta	503	104.0	44,656	9,235.4	5	1.0	1.0	483.5
Sweden	26,670	261.9	148,500	1,458.3	3,256	12.2	32.0	10,183.2
Netherlands	42,788	248.3	209,718	1,217.1	5,456	12.8	31.7	17,231
Republic of Korea	10,936	21.2	680,890	1,318.7	258	2.4	0.5	51,635.3

†Number of cases, number of deaths and population (in thousands) for all countries, but Cyprus, as reported by ECDC at

<https://www.ecdc.europa.eu/en/publications-data/download-todays-data-geographic-distribution-covid-19-cases-worldwide>

§ Data for Cyprus: internal communication; data for other countries: <https://www.finddx.org/covid-19/test-tracker/>

° CFR: Case fatality ratio.

\* Data from Statistical Service of the Republic of Cyprus, 2018 ([Statistical Service of the Republic of Cyprus](#))



## **Acknowledgments**

We thank Antry Constantinou, Christiana Soteriou, Despina Ioannou, Irene Georgiou, Maria Clerides, Tatiana Sofocleous, and Despoina Stylianou from the Ministry of Health, and the companies CELLOCK (<https://cellock.com>) and GEOMATIC (<https://geomatic.com>) for their volunteering support.



## Appendix

Table A1: Laboratory-confirmed COVID-19-cases in Cyprus by district of residence and origin (n = 904).

District/ <i>municipality</i>	Total		Travel-related		Unknown origin		Locally-acquired			Pop.
	N	%	N	%	N	%	N	%	N (per 100,000 pop)	
Ammochostos	41	4.5	9	7.0	9	6.3	23	3.6	47.7	48,200
Larnaka	234	25.9	18	14.0	47	32.6	169	26.8	115.0	147,000
<i>Aradippou</i>	123	13.6	10	7.8	33	22.9	80	12.7	416.1	19,228
Limassol	103	11.4	28	21.7	14	9.7	61	9.7	24.9	244,900
Nicosia	350	38.7	47	36.4	48	33.3	255	40.4	74.6	341,700
Pafos	155	17.1	11	8.5	25	17.4	119	18.9	126.5	94,100
Other	21	2.3	16	12.4	1	0.7	4	0.6		
<b>Total</b>	<b>904</b>	<b>100</b>	<b>129</b>	<b>100</b>	<b>144</b>	<b>100</b>	<b>631</b>	<b>100</b>	<b>72.0</b>	<b>875,900</b>

Other includes British Bases, abroad and unknown



Table A2: Sex and age distribution of asymptomatic cases at diagnosis (n = 280).

Characteristics	All cases (n = 904)	Asymptomatic cases (n = 280)	
	N	n	%
Sex			
Male	454	156	34.4
Female	450	124	27.6
Age groups (years)			
0-9	29	11	37.9
10-19	37	16	43.2
20-29	117	42	35.9
30-39	179	66	36.9
40-49	157	46	29.3
50-59	162	42	25.9
60-69	112	21	18.8
70-79	84	29	34.5
80+	27	7	25.9
Median age in years (IQR*)	45 (32-59)	40 (30-56)	

\*IQR: Interquartile Range



Table A3: Characteristics of all deaths (n = 23).

Characteristics	N	%
Sex		
Male	17	73.9
Female	6	26.1
Age groups (years)		
0-9	0	0.0
10-19	0	0.0
20-29	0	0.0
30-39	0	0.0
40-49	1	4.3
50-59	2	8.7
60-69	6	26.1
70-79	10	43.5
80+	4	17.4
Median age in years (IQR*)	76 (66-79)	
District		
Ammochostos	3	13.0
Larnaka	9	39.2
Limassol	2	8.7
Nicosia	3	13.0
Pafos	6	26.1

\*IQR: Interquartile Range





Table A4: Number of cases by date of sampling, laboratory reporting, death, and ICU admission.

Date	Sampling (n = 904)	Laboratory reporting (n = 904)	Death (n = 23)	ICU admission (n = 32)
01-Mar	0	0	0	0
02-Mar	0	0	0	0
03-Mar	0	0	0	0
04-Mar	0	0	0	0
05-Mar	0	0	0	0
06-Mar	0	0	0	0
07-Mar	1	0	0	0
08-Mar	0	0	0	0
09-Mar	1	2	0	0
10-Mar	4	0	0	0
11-Mar	2	0	0	0
12-Mar	6	0	0	0
13-Mar	12	16	0	0
14-Mar	8	5	0	0
15-Mar	12	8	0	0
16-Mar	5	13	0	1
17-Mar	8	4	0	1
18-Mar	16	16	0	0
19-Mar	13	9	0	0
20-Mar	17	10	0	1
21-Mar	10	0	1	1
22-Mar	6	16	0	1
23-Mar	13	19	0	1
24-Mar	18	8	2	3
25-Mar	14	10	0	3
26-Mar	34	20	0	1
27-Mar	31	24	3	2
28-Mar	26	21	1	3
29-Mar	33	27	1	1
30-Mar	37	33	0	0
31-Mar	39	45	2	0
01-Apr	29	56	2	1
02-Apr	47	29	0	0
03-Apr	21	32	1	2



**ΥΠΟΥΡΓΕΙΟ ΥΓΕΙΑΣ**

04-Apr	25	38	1	0
05-Apr	9	18	0	0
06-Apr	37	23	0	1
07-Apr	39	23	0	1
08-Apr	23	32	0	1
09-Apr	17	31	1	1
10-Apr	18	20	0	1
11-Apr	37	20	1	0
12-Apr	21	16	1	1
13-Apr	26	41	0	0
14-Apr	25	25	0	0
15-Apr	15	16	0	0
16-Apr	9	19	0	2
17-Apr	4	15	0	0
18-Apr	6	7	0	0
19-Apr	0	1	0	0
20-Apr	11	6	0	1
21-Apr	6	13	0	0
22-Apr	7	5	1	0
23-Apr	13	7	1	0
24-Apr	2	12	0	0
25-Apr	9	3	0	0
26-Apr	4	2	1	1
27-Apr	16	11	0	0
28-Apr	5	16	0	0
29-Apr	9	4	0	0
30-Apr	6	5	0	0
01-May	7	8	0	0
02-May	3	7	0	0
03-May	2	4	0	0
04-May	4	6	0	0
05-May	5	4	1	0
06-May	6	4	0	0
07-May	2	3	0	0
08-May	3	3	0	0
09-May	8	3	1	0
10-May	1	4	0	0
11-May	0	3	0	0



**ΥΠΟΥΡΓΕΙΟ ΥΓΕΙΑΣ**

12-May

1


3

1

0

---


Figure A1: Distribution of cases by postal code (n = 880 with information available).



Each colour represents a different postal code and the size changes according to the number of cases.



Figure A2: Cumulative tests and cases per 100,000 population in Cyprus and other selected countries (Updated: 12/05/2020).



Data source for Cyprus: internal communication; data source for other countries:

<https://www.finddx.org/covid-19/test-tracker/>

*Numbers of cases, tests and deaths for Cyprus are aggregated and include people from abroad and the British bases, while the total population does not include inhabitants from abroad or from the British bases.*



Figure A3: Time from date of sampling to death of COVID-19 cases who died (n = 23; for three cases who died on the day of sampling/reporting, the time alive has been considered 0.5 days).






Figure A4: Length of stay in ICU (n = 32; for two cases who died on the same day of ICU admission the length of stay in ICU has been considered 0.5 days).






Figure A5: Length of stay in ICU of patients who died and had been admitted to an ICU (n = 15; for two cases who died the same day of ICU admission the length of stay in ICU has been considered 0.5 days).

