



Coronavirus Disease 2019 (COVID-19)

National Surveillance Report as of 25/08/2020

Date of publication: 27/08/2020

Produced by the Epidemiological Surveillance Unit of the Ministry of Health

Contributors: Annalisa Quattrocchi, Ioannis Mamais, Valentinos Silvestros, Anna Demetriou (Health Monitoring Unit), Maria Athanasiadou (Health Monitoring Unit), Theopisti Kyprianou (Health Monitoring Unit), Androulla Stylianou, Sotiroula Sotiriou, Fani Theofhanous, Christos Charalambous, Ioanna Gregoriou, Maria Koliou, Georgios Nikolopoulos and Elisavet Constantinou

Scientific Committee: Elisavet Constantinou, Constantinos Constantinou, Niki Paphitou, Georgios Nikolopoulos, Maria Koliou, George Panos, Eirini Christaki, Zoi – Dorothea Pana, Constantinos Tsioutis, Leondios Kostrikis, Peter Karayiannis, George Petrikkos, Petros Agathangelou, George Mixides

Suggested citation: Epidemiological Surveillance Unit of the Ministry of Health, Cyprus. National Situation Report. Coronavirus Disease 2019 (COVID-19), 27 August 2020.



Summary

- As of August 25th, a total of 1,442 COVID-19 cases and 27 deaths (case fatality rate: 1.9%) have been reported in the Republic of Cyprus.
- Among these cases, 14.1% are health-care workers (n = 204) - 2.9% physicians (n = 42), 7% nurses (n = 101), 0.7% other health occupations (n = 10), and 3.5% auxiliary staff (n = 51).
- The median age of cases is 39 years (Interquartile range - IQR: 27-56 years); 51.4% are male and 48.6% are female.
- Overall, of 1,276 cases for which the place of exposure was known, locally acquired infections (index cases and close-contacts of confirmed cases) were 947 (74.2%) - of these 6% (n = 57) were related to a health-care facility (General Hospital in Pafos) and 13.5% (n = 128) were reported in Aradippou municipality.
- Since July 1st (included), of 444 cases reported, 34% (n = 153) were imported, 61% (n = 269) were locally-acquired, and 5% (n = 22) were unknown.
- In total, 14.1% (n = 203) of cases received hospital care, and six (3%) are still hospitalised (either for treatment of COVID-19 symptoms or for pre-existing conditions). The median age of all hospitalized patients is 62 years (IQR: 47-73 years) and 63.6% are males.
- There is one patient in intensive care unit.
- Among cases alive, 1,139 (80.5%) cases have recovered (without symptoms and with two negative tests following their diagnosis or released 21 days after diagnosis).
- A total of 285,300 tests have been performed as of August 25th (32,572.2 per 100,000 population).



Epidemiological surveillance in the Republic of Cyprus

Analyses are based on laboratory-confirmed cases notified to the Epidemiological Surveillance Unit of the Ministry of Health.

As of August 25th, 1,442 laboratory-confirmed cases of coronavirus disease 2019 (COVID-19) have been reported (Figure 1 and 2).

The median time between symptoms onset and date of sampling was 3 days (Interquartile range - IQR: 2-6 days). It should be noted that for 32 cases the date of sample collection was before the onset of symptoms because of immediate testing of contacts of possible and laboratory-confirmed cases.

As of August 25th, the 14-day cumulative incidence rate of COVID-19 (per 100,000 population), a measure which reflects the number of active COVID-19 cases in the population (prevalence)¹, is 18.4 per 100,000 population (Figure 3).




Figure 1: Number of laboratory-confirmed COVID-19 cases in Cyprus since 03/03/2020 by date of sample collection and date of symptoms onset (n = 1,440 and n = 706 with data available, respectively).

Recent data should be interpreted with caution due to the possibility that cases with date of onset within the reporting period have not yet been diagnosed.

¹Coronavirus disease 2019 (COVID-19) pandemic: increased transmission in the EU/EEA and the UK – seventh update, 25 March 2020. Stockholm: ECDC; 2020.

<https://www.ecdc.europa.eu/sites/default/files/documents/RRA-seventh-update-Outbreak-of-coronavirus-disease-COVID-19.pdf>




Figure 2: Number and cumulative number of laboratory-confirmed COVID-19 cases in Cyprus since 03/03/2020, by date of laboratory reporting (n = 1,442).
Recent data should be interpreted with caution due to the possibility that cases with date of onset within the reporting period have not yet been diagnosed.




Figure 3. COVID-19 14-day cumulative incidence rate per 100,000 population (proxy of COVID-19 prevalence).
March 23rd represents the first 14th day since cases have been reported.

Characteristics of the cases

Among these cases, 51.4% are male (n = 741) and 48.6% female (n = 701). The median age of cases is 39 years (IQR: 27-56 years). By age group, cases included 106 infants, children and adolescents aged 0-17 years-old (7.4%), 1,063 adults aged 18-59 years (73.7%), and 273 persons aged 60 years and older (18.9%) (Figure 4).




Figure 4: Laboratory-confirmed COVID-19 cases in Cyprus by sex and age groups.

Among all cases, 476 (33%) were reported in Nicosia district, 372 (25.8%) in Larnaka, 256 (17.8%) in Limassol, 183 (12.7%) in Pafos, 74 (5.1%) in Ammochostos, and 81 (5.6%) were reported either in the British bases or had a residence abroad, or information was not available (Table A1 in appendix).

Figure A1 in appendix shows the distribution of cases by postal code.

Notably, 176 cases (12.2%) were reported in Aradippou, a municipality in Larnaka district (Table A1 in appendix). Cases in Aradippou, including a cluster in a local bakery production line, are mainly males (51.7%; n = 91) and the median age is 39 years (IQR: 26-57 years). If the cluster is excluded, cases are mainly females (57.4%; n = 85) and the median age is 45 years (IQR: 24-60 years).



Among the 1,442 cases, 14.1% are health-care workers² (n = 204) - 2.9% physicians (n = 42), 7% nurses (n = 101), 0.7% other health occupations (n = 10), and 3.5% auxiliary staff (n = 51). Table 1 shows the distribution of health-care workers by district of residence.

Table 1: Health-care workers by district of residence (n=204).

District	Health-care worker	Physicians	Nurses	Other health occupations	Auxiliary staff
Ammochostos	16	3	7	0	6
Larnaka	46	8	25	2	11
Limassol	22	3	11	2	6
Nicosia	60	13	26	4	17
Pafos	60	15	32	2	11
Total	204	42	101	10	51

Epidemiological link

As of August 25th, place of exposure is available for 1,276 cases (88.5%). In total, 25.8% (n = 329) of laboratory-confirmed COVID-19-cases had history of travel or residence abroad during the 14 days prior to symptom onset (imported). Locally-acquired infections (index cases and close-contacts of confirmed cases) occurred in 74.2% (n = 947 of 1,276 with known place of exposure) of the cases, of which 6% (n = 57) were related to a health-care facility (General Hospital in Pafos). Of all cases in Aradippou (Larnaka district) (n = 176), 128 (72.7%) were locally-acquired, 13 (7.4%) imported and for 35 cases (19.9%) the epidemiological link was not recorded. Table A1 in the appendix shows the number and the rate (per 100,000 population) of locally-acquired cases by district of residence.

Since July 1st (included), of 444 cases reported, 34% (n = 153) were imported, 61% (n = 269) were locally-acquired and 5% (n = 22) are unknown. Table B1 and Figure B1 in Appendix show the characteristics of cases by origin of infection and the number of cases by origin of infection in the recent period, respectively. Figure B2 shows the distribution of cases by origin of infection for each district. The majority of recent cases were reported in Limassol (31.5%; n = 133), where 70.7% of them (n = 94) were locally-acquired.

² The term “health-care worker” is based on the occupation and not on the place of exposure. Health-care workers are defined as all health care professionals, allied health workers, and auxiliary health workers.



Clinical features

Of the 1,442 laboratory-confirmed COVID-19-cases, clinical information is available for 97.9% (n = 1,411), of which 38.9% (n = 549) reported no symptoms at diagnosis and 61.1% (n = 862) reported at least one symptom. The most commonly reported symptoms were:

- cough (381/1,391; 27.4%),
- fever (397/1,393; 28.5%),
- myalgia (264/1,389; 19%),
- sore throat (221/1,388; 15.9%),
- anosmia (158/1,300; 12.2%), and
- shortness of breath (132/1,272; 9.6%).

Other reported symptoms were diarrhoea, runny nose, and headache.

Table A2 in appendix reports the sex and age distribution of asymptomatic cases at diagnosis.

Pre-existing conditions

Information on comorbidities was available for 1,307 (90.6%) cases. Of these, 448 (34.2%) reported at least one comorbidity.

The most commonly reported comorbidities were:

- hypertension (158/1,295; 12.2%),
- diabetes (88/1,300; 6.7%),
- heart disease (77/1,300; 5.9%), and
- cancer (29/1,307; 2.2%).

Other reported comorbidities were immunosuppression, chronic respiratory disease, chronic kidney disease, and autoimmune disease.

Deaths

As of August 25th, 27 deaths were reported in Cyprus (Case Fatality Rate - CFR: 1.9%).

The mortality rate for COVID-19 is 3.1 per 100,000 population.

Twenty deaths (74%) occurred in men and seven (26%) in women; the median age of all deaths was 76 years (IQR: 66-79 years). Ten deaths were reported among residents in Larnaka, eight in Pafos, four in Nicosia, three in Ammochostos, and two in Limassol (Appendix Table A3).

The median time from date of sampling to death was 12 days (IQR: 4-30 days). Figure A3 shows the Kaplan-Meier curve of the time from date of sampling to death.

For 20 deaths, COVID-19 was the underlying cause of death (COVID-19 CFR: 1.4%).

Figure 5 reports the number of deaths by date.




Figure 5: Number of deaths among COVID-19 cases in Cyprus by date of death (n = 27).

Hospitalization and intensive care unit (ICU) admissions³

In total, 14.1% (n = 203) of people with COVID-19 received hospital care, and six (3%) are still hospitalised (either for treatment of COVID-19 symptoms or for pre-existing conditions). The median age of hospitalized patients was 62 years (IQR: 47-73 years). Hospitalized cases were mainly males (n = 129; 63.6%).

Figure 6 shows the total number of first hospital admissions by date.




Figure 6: Number of laboratory-confirmed COVID-19 cases by date of first hospital admission (n = 203).

**Date of hospital admission was replaced with date of sampling for inpatients hospitalised prior to the beginning of the epidemic.*

Overall, 33 cases (16.3% of all hospitalized patients) have been admitted to ICU⁴, and currently there is one case in ICU (as of August 25th).

A total of 28 ICU patients (84.8% of all ICU patients) have been intubated, and currently there is one patient intubated.

The overall median length of stay in ICU (for all 33 ICU cases) was 11 days (IQR: 8-28 days). Figure A4 shows the Kaplan-Meier curve of the length of stay in ICU.

For patients who died while in ICU (n = 18), the median length of stay in ICU was 13.5 days (IQR: 8-28). Figure A5 shows the Kaplan-Meier curve of the length of stay in ICU for the people who died.

³ Data on hospitalisation and ICU are provisional and should be interpreted with caution because delay in data reporting is likely; for the construction of the curve, people are no longer in an ICU the day next to the date of their discharge, death or transfer.

⁴ Intensive care unit (ICU) refers only to the ICU in Limassol General Hospital and to the ICU in Nicosia General Hospital.



For patients transferred/discharged alive from ICU (n = 14), the median length of stay in ICU was 10.5 days (IQR: 8-28 days).

The median age of patients ever admitted to ICU was 66 years (IQR: 56-76 years). ICU patients are mainly male (n = 24; 72.7%).

The characteristics of patients ever admitted to ICU and discharged or who died in ICU are reported in Table B2 in Appendix.

The number of cases currently in ICU is 0.1 per 100,000 population.

Figure 7 shows the number of patients in ICU, by day and intubation. Table A4 in the appendix shows the total number of ICU admissions by date.




Figure 7: Number of laboratory-confirmed COVID-19 cases in ICU by date and intubation.

**Date of discharge/transfer/death included*

Recovered/released

As of August 25th, among cases alive, 80.5% (n = 1,139) of COVID-19 cases have recovered⁵; of which 914 (80.3%) tested negative two consecutive times, and 225 (19.7%) have been released as per the new guidelines⁶. The median time between the second negative result and the first date of sampling was 23 days (IQR: 18-36 days).

Table 2 shows the number and percentage of recovered cases and their characteristics.

Table 2: Characteristics of all cases and cases recovered/released (n = 1,139)

Characteristics	Total cases N	Recovered			
		Two consecutive negative tests		Released after 21 days	
		N	%	N	%
Total	1,442	914	63.4	225	15.6
Sex					
Male	741	462	62.3	130	17.5
Female	701	452	64.5	95	13.6
Age groups (years)					
0-9	46	22	47.8	11	23.9
10-19	106	50	47.2	25	23.6
20-29	289	155	53.6	53	18.3
30-39	284	184	64.8	41	14.4
40-49	220	151	68.6	37	16.8
50-59	224	168	75.0	23	10.3
60-69	142	99	69.7	20	14.1
70-79	98	66	67.3	12	12.2
80+	33	19	57.6	3	9.1
Median age in years (IQR*)	39 (27-56)	43 (30-57)		35 (24-50)	

⁵ For symptomatic cases, or for cases hospitalised, a COVID-19 case can be considered cured after the resolution of symptoms and two negative tests for SARS-CoV-2 at 24-hour interval at least. For asymptomatic cases, or for persons isolated at home, the negative tests to document virus clearance should be obtained at a minimum of 14 days after the initial positive test (end of the quarantine period).

Novel coronavirus (SARS-CoV-2). Discharge criteria for confirmed COVID-19 cases- When is it safe to discharge COVID-19 cases from the hospital or end home isolation? - Technical Report, 10 March 2020. Stockholm: ECDC; 2020.

⁶ A person is released 21 days after the initial diagnosis, if he/she has a positive test 14 days after the initial diagnosis and remains in isolation for one more week without being further tested.



Comparison with selected countries

As of August 25th, in Cyprus the reporting rate was 164.6 cases per 100,000 population, the mortality rate was 3.1 deaths per 100,000 population and the CFR was 1.9%.

Table 3 shows COVID-19 indicators for Cyprus and other selected countries.

Figure A2 in appendix reports the rates of cumulative tests and cases (per 100,000 population) in Cyprus and other selected countries. In Cyprus the testing rate is 32,572.2 per 100,000 population.

It should be noted that the number of cases, tests and deaths for Cyprus are aggregated and include people from abroad and the British bases, while the total population does not include inhabitants from abroad or from the British bases.



Table 3: COVID-19 indicators by selected countries, as of 25/08/2020

Country	N. of cases [†]	N. of cases (per 100,000 pop)	N. of tests [§]	N. of tests (per 100,000 pop)	N. of deaths [†]	CFR [°] (%)	Mortality rate (per 100,000 pop)	Pop. (in thousand) [†]
Cyprus	1,442	164.6	285,300	32,572.2	27	1.9	3.1	875.9*
Italy	260,298	431.2	8,125,892	13,462.5	35,441	13.6	58.7	60,359.5
USA	5,740,909	1,744.6	72,382,318	21,996.4	177,279	3.1	53.9	329,064.9
UK	326,614	490.1	15,177,265	22,772.6	41,433	12.7	62.2	66,647.1
Greece	8,819	82.2	858,138	8,001.6	242	2.7	2.3	10,724.6
Malta	1,667	337.8	177,518	35,966.9	10	0.6	2.0	493.6
Sweden	86,721	847.7	783,467	7,658.4	5,813	6.7	56.8	10,230.2
Netherlands	67,062	388.0	411,972	2,383.8	6,193	9.2	35.8	17,282.2
Republic of Korea	17,945	35.0	1,825,837	3,564.3	310	1.7	0.6	51,225.3

[†]Number of cases, number of deaths and population (in thousands) for all countries, but Cyprus, as reported by ECDC at <https://www.ecdc.europa.eu/en/publications-data/download-todays-data-geographic-distribution-covid-19-cases-worldwide>

[§] Data for Cyprus: internal communication; data for other countries: <https://www.finndx.org/covid-19/test-tracker/>

[°] CFR: Case fatality ratio.

* Data from Statistical Service of the Republic of Cyprus, 2018 ([Statistical Service of the Republic of Cyprus](#))



Acknowledgments

We thank Antry Constantinou, Christiana Soteriou, Despina Ioannou, Irene Georgiou, Maria Clerides, Tatiana Sofocleous, and Despoina Stylianou from the Ministry of Health, and the companies CELLOCK (<https://cellock.com>) and GEOMATIC (<https://geomatic.com>) for their volunteering support.



Appendix

Table A1: Laboratory-confirmed COVID-19-cases in Cyprus by district of residence and origin (n = 1,442).

District/ <i>municipality</i>	Total		Travel-related		Unknown origin		Locally-acquired			Pop.
	N	%	N	%	N	%	N	%	N (per 100,000 pop)	
Ammochostos	74	5.1	25	7.6	9	5.4	40	4.2	83.0	48,200
Larnaka	372	25.8	38	11.6	56	33.7	278	29.4	189.1	147,000
<i>Aradippou</i>	176	12.2	13	4.0	35	21.1	128	13.5	665.7	19,228
Limassol	256	17.8	79	24.0	18	10.8	159	16.8	64.9	244,900
Nicosia	476	33.0	97	29.5	54	32.5	325	34.3	95.1	341,700
Pafos	183	12.7	22	6.7	27	16.3	134	14.1	142.4	94,100
Other	81	5.6	68	20.7	2	1.2	11	1.2		
Total	1442	100	329	100	166	100	947	100	108.1	875,900

Other includes British Bases, abroad and unknown



Table A2: Sex and age distribution of asymptomatic cases at diagnosis (n = 549).

Characteristics	All cases (n = 1,442)	Asymptomatic cases (n = 549)	
	N	n	%
Sex			
Male	741	307	41.4
Female	701	242	34.5
Age groups (years)			
0-9	46	22	47.8
10-19	106	46	43.4
20-29	289	136	47.1
30-39	284	123	43.3
40-49	220	72	32.7
50-59	224	72	32.1
60-69	142	36	25.4
70-79	98	32	32.7
80+	33	10	30.3
Median age in years (IQR*)	39 (27-56)	35 (25-51)	

*IQR: Interquartile Range

Table A3: Characteristics of all deaths (n = 27) and due to COVID-19 (n = 20)

Characteristics	All deaths		COVID-19 deaths	
	N	%	N	%
Male	20	74.1	14	70.0
Female	7	25.9	6	30.0
Age groups (years)				
0-9	0	0.0	0	0.0
10-19	0	0.0	0	0.0
20-29	0	0.0	0	0.0
30-39	0	0.0	0	0.0
40-49	1	3.7	1	5.0
50-59	2	7.4	1	5.0
60-69	8	29.6	7	35.0
70-79	11	40.7	9	45.0
80+	5	18.5	2	10.0
Median age in years (IQR*)	76 (66-79)		75 (65.5-77.5)	
District				0.0
Ammochostos	3	11.1	1	5.0
Larnaka	10	37.0	7	35.0
Limassol	2	7.4	2	10.0
Nicosia	4	14.8	4	20.0
Pafos	8	29.6	6	30.0
Comorbidities				0.0
No	3	11.1	11	55.0
Yes	23	85.2	8	40.0
Median age in years (IQR*)	74 (66-79)		72.5 (67.5-76.5)	
Unknown	1	3.7	1	5.0
Diabetes	10	37.0	8	40.0
Median age in years (IQR*)	74 (69-77)		72.5 (67.5-76.5)	
Hypertension	8	29.6	6	30.0
Median age in years (IQR*)	76 (70.5-81.5)		74 (69-77)	
Heart disease	12	44.4	10	50.0
Median age in years (IQR*)	75 (67.5-80)		75 (66-77)	
Chronic kidney disease	6	22.2	3	15.0
Median age in years (IQR*)	78 (69-83)		76 (63-85)	
Chronic respiratory disease	4	14.8	4	20.0
Median age in years (IQR*)	75 (71-77)		75 (71-77)	



ΥΠΟΥΡΓΕΙΟ ΥΓΕΙΑΣ

Chronic liver disease	3	11.1	2	10.0
Median age in years (IQR*)	76 (69-79)		77.5 (76-79)	
Immunosuppression	0	0.0	0	0.0
Median age in years (IQR*)				
Cancer	4	14.8	1	5.0
Median age in years (IQR*)	81 (67.5-88)		79	
Autoimmunedisease	0	0.0	0	0.0
Median age in years (IQR*)				

*IQR: Interquartile Range



ΥΠΟΥΡΓΕΙΟ ΥΓΕΙΑΣ

Table A4: Number of cases by date of sampling, laboratory reporting, death, and ICU admission.

Date	Sampling (n = 1,440)	Laboratory reporting (n = 1,442)	Death (n = 27)	ICU first admission (n = 33)
01-Mar	0	0	0	0
02-Mar	0	0	0	0
03-Mar	0	0	0	0
04-Mar	0	0	0	0
05-Mar	0	0	0	0
06-Mar	0	0	0	0
07-Mar	1	0	0	0
08-Mar	0	0	0	0
09-Mar	1	2	0	0
10-Mar	4	0	0	0
11-Mar	2	0	0	0
12-Mar	6	0	0	0
13-Mar	13	16	0	0
14-Mar	8	5	0	0
15-Mar	12	8	0	0
16-Mar	5	13	0	1
17-Mar	8	4	0	1
18-Mar	16	16	0	0
19-Mar	13	9	0	0
20-Mar	16	10	0	1
21-Mar	10	0	1	1
22-Mar	6	16	0	1
23-Mar	13	19	0	1
24-Mar	18	8	2	3
25-Mar	14	10	0	3
26-Mar	34	20	0	1
27-Mar	31	24	3	2
28-Mar	26	21	1	3
29-Mar	33	27	1	1
30-Mar	37	33	0	0
31-Mar	39	45	2	0
01-Apr	29	56	2	1
02-Apr	47	29	0	0

Data are subject to change due to the rapidly evolving situation



ΥΠΟΥΡΓΕΙΟ ΥΓΕΙΑΣ

03-Apr	21	32	1	2
04-Apr	25	38	1	0
05-Apr	9	18	0	0
06-Apr	37	23	0	1
07-Apr	39	23	0	1
08-Apr	23	32	0	1
09-Apr	17	31	1	1
10-Apr	18	20	0	1
11-Apr	37	20	1	0
12-Apr	21	16	1	1
13-Apr	26	41	0	0
14-Apr	25	25	0	0
15-Apr	15	16	0	0
16-Apr	9	19	0	2
17-Apr	4	15	0	0
18-Apr	6	7	0	0
19-Apr	0	1	0	0
20-Apr	11	6	0	1
21-Apr	6	13	0	0
22-Apr	7	5	1	0
23-Apr	13	7	1	0
24-Apr	2	12	0	0
25-Apr	9	3	0	0
26-Apr	4	2	1	1
27-Apr	16	11	0	0
28-Apr	5	16	0	0
29-Apr	9	4	0	0
30-Apr	6	5	0	0
01-May	7	8	0	0
02-May	3	7	0	0
03-May	2	4	0	0
04-May	4	6	0	0
05-May	5	4	1	0
06-May	6	4	0	0
07-May	2	3	0	0
08-May	3	3	0	0
09-May	8	3	1	0
10-May	1	4	0	0

Data are subject to change due to the rapidly evolving situation



ΥΠΟΥΡΓΕΙΟ ΥΓΕΙΑΣ

11-May	0	3	0	0
12-May	3	4	1	0
13-May	2	0	1	0
14-May	3	3	0	0
15-May	2	4	0	0
16-May	4	4	0	0
17-May	0	0	0	0
18-May	1	1	0	0
19-May	5	5	0	0
20-May	3	0	0	0
21-May	8	7	0	0
22-May	1	1	0	0
23-May	2	3	0	0
24-May	0	3	0	0
25-May	2	2	0	0
26-May	0	1	0	0
27-May	3	1	0	0
28-May	0	1	0	0
29-May	2	1	0	0
30-May	3	1	0	0
31-May	1	4	0	0
01-Jun	4	2	0	0
02-Jun	6	5	0	0
03-Jun	1	5	0	0
04-Jun	0	0	0	0
05-Jun	5	1	1	0
06-Jun	5	1	0	0
07-Jun	1	8	0	0
08-Jun	2	1	0	0
09-Jun	2	2	0	0
10-Jun	2	3	0	0
11-Jun	4	1	0	0
12-Jun	0	4	0	0
13-Jun	3	0	0	0
14-Jun	2	3	0	0
15-Jun	0	2	0	0
16-Jun	0	0	0	0
17-Jun	1	0	0	0



ΥΠΟΥΡΓΕΙΟ ΥΓΕΙΑΣ

18-Jun	0	0	1	0
19-Jun	1	1	0	0
20-Jun	1	0	0	0
21-Jun	0	1	0	0
22-Jun	1	1	0	0
23-Jun	3	2	0	0
24-Jun	0	1	0	0
25-Jun	2	1	0	0
26-Jun	0	2	0	0
27-Jun	0	0	0	0
28-Jun	2	0	0	0
29-Jun	2	4	0	0
30-Jun	1	0	0	0
01-Jul	0	1	0	0
02-Jul	1	0	0	0
03-Jul	1	0	0	0
04-Jul	3	4	0	0
05-Jul	0	1	0	0
06-Jul	2	0	0	0
07-Jul	2	3	0	0
08-Jul	2	3	0	0
09-Jul	3	2	0	0
10-Jul	8	1	0	0
11-Jul	0	8	0	0
12-Jul	1	0	0	0
13-Jul	1	1	0	0
14-Jul	2	2	0	0
15-Jul	7	5	0	0
16-Jul	0	3	0	0
17-Jul	5	5	0	0
18-Jul	1	1	0	0
19-Jul	0	0	0	0
20-Jul	2	1	0	0
21-Jul	0	1	0	0
22-Jul	5	4	0	0
23-Jul	2	3	0	0
24-Jul	9	7	0	0
25-Jul	3	2	0	0

Data are subject to change due to the rapidly evolving situation



ΥΠΟΥΡΓΕΙΟ ΥΓΕΙΑΣ

26-Jul	1	2	0	0
27-Jul	3	3	0	0
28-Jul	18	9	0	0
29-Jul	9	17	0	0
30-Jul	32	20	0	0
31-Jul	20	14	0	0
01-Aug	14	29	0	0
02-Aug	12	5	1	0
03-Aug	18	24	0	0
04-Aug	14	12	0	0
05-Aug	19	12	0	0
06-Aug	8	17	0	0
07-Aug	11	6	0	0
08-Aug	6	14	0	0
09-Aug	7	4	0	0
10-Aug	27	20	0	0
11-Aug	21	17	0	0
12-Aug	10	23	0	0
13-Aug	19	8	0	0
14-Aug	14	16	0	0
15-Aug	5	9	0	0
16-Aug	7	10	0	0
17-Aug	18	9	0	1
18-Aug	14	19	0	0
19-Aug	10	18	0	0
20-Aug	11	5	0	0
21-Aug	10	11	0	0
22-Aug	1	10	0	0
23-Aug	10	6	0	0
24-Aug	12	11	0	0
25-Aug	0	6	0	0




Figure A1: Distribution of cases by postal code (n = 1,340 with information available).



Each colour represents a different postal code and the size changes according to the number of cases.



Figure A2: Cumulative tests and cases per 100,000 population in Cyprus and other selected countries (Updated: 25/08/2020).



Data source for Cyprus: internal communication; data source for other countries:

<https://www.finndx.org/covid-19/test-tracker/>

Numbers of cases and tests for Cyprus are aggregated and include people from abroad and the British bases, while the total population does not include inhabitants from abroad or from the British bases.



Figure A3: Time from date of sampling to death of COVID-19 cases who died (n = 27; for three cases who died on the day of sampling/reporting, the time alive has been considered 0.5 days).






Figure A4: Length of stay in ICU (n = 33; for two cases who died on the same day of ICU admission the length of stay in ICU has been considered 0.5 days).







Figure A5: Length of stay in ICU of patients who died and had been admitted to an ICU (n = 18; for two cases who died the same day of ICU admission the length of stay in ICU has been considered 0.5 days).





ΥΠΟΥΡΓΕΙΟ ΥΓΕΙΑΣ

Table B1: Characteristics of cases by origin of infection, since July 1st


Characteristics	Total (n = 444)		Imported (n = 153)		Locally-acquired (n = 269)		Unknown (n = 22)	
	N	%	N	%	N	%	N	%
Sex								
Male	233	52.5	96	62.7	125	46.5	12	54.5
Female	211	47.5	57	37.3	144	53.5	10	45.5
Age groups (years)								
0-9	15	3.4	1	0.7	12	4.5	2	9.1
10-19	57	12.8	14	9.2	39	14.5	4	18.2
20-29	138	31.1	68	44.4	63	23.4	7	31.8
30-39	88	19.8	37	24.2	48	17.8	3	13.6
40-49	55	12.4	13	8.5	42	15.6	0	0.0
50-59	50	11.3	12	7.8	36	13.4	2	9.1
60-69	27	6.1	4	2.6	22	8.2	1	4.5
70-79	12	2.7	4	2.6	6	2.2	2	9.1
80+	2	0.5	0	0.0	1	0.4	1	4.5
Median age in years (IQR*)	30.5 (23-46)		29 (23-37)		33 (23-49)		26.5 (19-55)	
District								
Ammochostos	24	5.4	10	6.5	14	5.2	0	0.0
Larnaka	124	27.9	17	11.1	99	36.8	8	36.4
Limassol	137	30.9	39	25.5	94	34.9	4	18.2
Nicosia	91	20.5	34	22.2	50	18.6	7	31.8
Pafos	17	3.8	6	3.9	9	3.3	2	9.1
Other	51	11.5	47	30.7	3	1.1	1	4.5
Symptoms at diagnosis								
Yes	217	48.9	45	29.4	163	60.6	9	40.9
No	210	47.3	93	60.8	104	38.7	13	59.1
Unknown	17	3.8	15	9.8	2	0.7	0	0.0

Table B2. Characteristics of patients ever admitted to ICU, by patient status at ICU discharge

Characteristics	Total cases		Discharged alive		Deceased in ICU	
	N	%	N	%	N	%
Total	32	100.0	14	43.8	18	56.3
Sex						
Male	23	71.9	10	71.4	13	72.2
Female	9	28.1	4	28.6	5	27.8
Age groups (years)						
0-9	0	0.0	0	0.0	0	0.0
10-19	0	0.0	0	0.0	0	0.0
20-29	1	3.1	1	7.1	0	0.0
30-39	2	6.3	2	14.3	0	0.0
40-49	2	6.3	1	7.1	1	5.6
50-59	7	21.9	5	35.7	2	11.1
60-69	10	31.3	3	21.4	7	38.9
70-79	9	28.1	2	14.3	7	38.9
80+	1	3.1	0	0.0	1	5.6
Median age (IQR)	65.5 (56-75)		57 (47-69)		69 (65-76)	
Symptoms at diagnosis (yes)	29	90.6	13	92.9	16	88.9
Cough	16	50.0	7	50.0	9	50.0
Fever	16	50.0	8	57.1	8	44.4
Myalgia	4	12.5	2	14.3	2	11.1
Sore throat	4	12.5	2	14.3	2	11.1
Anosmia	0	0.0	0	0.0	0	0.0
Shortness of breath	10	31.3	3	21.4	7	38.9
Comorbidities (yes)	23	71.9	8	57.1	15	83.3
Diabetes	9	28.1	2	14.3	7	38.9
Hypertension	11	34.4	5	35.7	6	33.3
Heart diseases	9	28.1	1	7.1	8	44.4
Chronic kidney disease	3	9.4	0	0.0	3	16.7
Chronic respiratory disease	4	12.5	0	0.0	4	22.2
Chronic liver disease	1	3.1	0	0.0	1	5.6
Cancer	2	6.3	1	7.1	1	5.6
Immunosuppression	1	3.1	1	7.1	0	0.0



Figure B1. Number of cases by origin of infection, since July 1st





ΥΠΟΥΡΓΕΙΟ ΥΓΕΙΑΣ

Figure B2. Number of cases by origin of infection for each district, since July 1st (n = 444)

