



## **Coronavirus Disease 2019 (COVID-19)**

### **National Surveillance Report as of 09/02/2021**

Date of publication: 12/02/2021

Produced by the Epidemiological Surveillance Unit of the Ministry of Health

Contributors: Annalisa Quattrocchi, Ioannis Mamais, Valentinos Silvestros, Anna Demetriou (Health Monitoring Unit), Maria Athanasiadou (Health Monitoring Unit), Theopisti Kyprianou (Health Monitoring Unit), Androulla Stylianou, Sotiroula Sotiriou, Fani Theofhanous, Christos Charalambous, Ioanna Gregoriou, Maria Koliou, Olga Kalakouta, Georgios Nikolopoulos

Scientific Committee: Constantinos Constantinou, Niki Paphitou, Georgios Nikolopoulos, Maria Koliou, George Panos, Eirini Christaki, Zoi – Dorothea Pana, Constantinos Tsioutis, Peter Karayiannis, George Petrikkos, Petros Agathangelou, George Mixides

Suggested citation: Epidemiological Surveillance Unit of the Ministry of Health, Cyprus. National Situation Report. Coronavirus Disease 2019 (COVID-19), 12 February 2021.



## Summary

- As of February 9<sup>th</sup>, 2021, a total of 31,704 COVID-19 cases have been diagnosed, of which 217 died due to COVID-19 (case fatality risk: 0.7%).
- In the last 14 days (27 Jan - 09 Feb, 2021), 1,536 cases were diagnosed. The 14-day cumulative diagnosis rate is 173 per 100,000 population.
  - The median age was 38 years (Interquartile range - IQR: 25-54 years); 50.3% were female (n = 773), 49.6% male (n = 762), and for one case the information is not currently available,
  - By place of exposure: 1% (n = 16) were imported and 99% (n = 1,520) were locally-acquired.
- As of February 10<sup>th</sup>, 2021, 98 people were still hospitalized (currently notified). The median age of patients still hospitalized as of 10<sup>th</sup> of February, 2021 (n = 98, including those discharged on that day) is 67 years (IQR: 53-75 years), 52% (n = 51) are males, and 44.9% (n = 44) are from Limassol district.
- Of 23 cases in intensive care units (ICU), who are currently notified, all are intubated. The median age of current ICU patients is 70 (IQR: 60-76) years and 13 (56.5%) are females. Twelve (52.2%) patients currently in ICU have pre-existing conditions.
- Over the last 14 days, 25,665 RT PCR and 362,452 rapid antigen tests have been performed (2,890.2 RT PCR and 40,816.7 rapid antigen tests per 100,000 population).



## Epidemiological surveillance in the Republic of Cyprus

Analyses are based on laboratory-confirmed cases notified to the Epidemiological Surveillance Unit of the Ministry of Health.

As of February 9<sup>th</sup>, 2021, 31,704 cases of coronavirus disease 2019 (COVID-19) have been diagnosed (laboratory-confirmed) (Figure 1, 2 and 3).

In the last 14 days (27 Jan-09 Feb, 2021), 1,536 cases have been diagnosed. The 14-day cumulative diagnosis rate of COVID-19 (per 100,000 population)<sup>1</sup>, a measure which reflects the number of active COVID-19 cases in the population (prevalence)<sup>2</sup>, is 173 per 100,000 population (Figure 3).




Figure 1: Number and 7-day moving average (last moving average on Feb 9<sup>th</sup>, 2021, is shown on the graph) of laboratory-confirmed COVID-19 cases in the Republic of Cyprus since 03/03/2020 by date of sample collection\*.

\* when date of sample collection was not available, laboratory result date or hospitalisation date was used.

<sup>1</sup>Population denominator for end of year 2019, available at [https://www.mof.gov.cy/mof/cystat/statistics.nsf/populationcondition\\_21main\\_en/populationcondition\\_21main\\_en?OpenForm&sub=1&sel=4](https://www.mof.gov.cy/mof/cystat/statistics.nsf/populationcondition_21main_en/populationcondition_21main_en?OpenForm&sub=1&sel=4)

<sup>2</sup>Coronavirus disease 2019 (COVID-19) pandemic: increased transmission in the EU/EEA and the UK – seventh update, 25 March 2020. Stockholm: ECDC; 2020. <https://www.ecdc.europa.eu/sites/default/files/documents/RRA-seventh-update-Outbreak-of-coronavirus-disease-COVID-19.pdf>




Figure 2: Number of laboratory-confirmed COVID-19 cases in the Republic of Cyprus since 03/03/2020 by date of symptoms onset.

**Recent data should be interpreted with caution due to the possibility that cases with date of onset within the reporting period have not yet been diagnosed.**




Figure 3: COVID-19 14-day cumulative diagnosis rate per 100,000 population (proxy of COVID-19 prevalence).



## Characteristics of cases diagnosed in the last 14 days

### Age, sex, and place of residence

Among 1,536 cases diagnosed since 27<sup>th</sup> January, 2021, 50.3% were female (n = 773), 49.6% male (n = 762), and for one case the information is not currently available.

By age group, cases included 253 infants, children and adolescents aged 0-19 years-old (16.5%), 1,018 adults aged 20-59 years (66.3%), and 265 persons aged 60 years and older (17.2%). Figure 4 shows the number of cases by 10-year age band and sex. The median age of all cases diagnosed in the last 14 days is 38 years (IQR: 25-54 years). The median age of adult cases ( $\geq 18$  years) is 42 years (IQR: 30-56 years). The 14-day cumulative diagnosis rate (per 100,000 population) by age group is shown in figure A1 in the Appendix.




Figure 4: Laboratory-confirmed COVID-19 cases in the Republic of Cyprus, diagnosed in the last 14 days, by age group and sex.

Among all cases diagnosed in the last 14 days, 835 (54.4%) were reported in Limassol district, 370 (24.1%) in Nicosia, 181 (11.8%) in Larnaka, 61 (4%) in Pafos, 51 (3.3%) in Ammochostos, and 38 (2.5%) were reported either in the British bases or had a residence abroad, or information was not available. Figure A2 in the Appendix shows the 14-day cumulative diagnosis rate (per 100,000 population) by district.



### Epidemiological link

In the last 14 days (27 Jan-09 Feb, 2021), of 1,536 cases sampled and notified, 1% (n = 16) were imported and 99% (n = 1,520) were locally-acquired. Table A1 and Figure A3 (A&B) in the Appendix show the characteristics of cases by origin of infection and the daily number of locally-acquired cases by district in the recent period, respectively.

### Clinical features

In the last 14 days (27 Jan-09 Feb, 2021), 35.4% (n = 644) reported no symptoms at diagnosis and 64.6% (n = 992) reported at least one symptom.

### Pre-existing conditions

In the last 14 days (27 Jan-09Feb, 2021), 375 cases (24.4%) reported at least one comorbidity.



## Deaths

Among cases, 217 COVID-19 associated deaths were reported in the Republic of Cyprus until February 10<sup>th</sup>, 2021 (Case Fatality Risk - CFR: 0.7%).

The COVID-19 associated mortality is 24.4 per 100,000 population.

Deaths occurred in 148 men (68.2%) and 69 (31.8%) women; the median age of all COVID-19 associated deaths was 81 years (IQR: 73-87 years). By district of residence, deceased cases were 74 (34.1%) from Nicosia, 64 (29.5%) from Limassol, 42 (19.4%) from Larnaka, 17 (7.8%) from Pafos, 14 (6.5%) from Ammochostos and six deaths (2.8%) occurred among cases reported either in the British bases or had a residence abroad, or information was not available.

The median time from date of sampling to death (due to COVID-19) was 13 days (IQR: 7-20 days).

Figure 5a reports the number of COVID-19 associated deaths by date.

Among cases, 264 deaths (all causes) were reported in the Republic of Cyprus until February 10<sup>th</sup>, 2021 (CFR: 0.8%).

The mortality (all-causes) for people with COVID-19 is 29.7 per 100,000 population.

Deaths occurred in 183 men (69.3%) and 81 (30.7%) women; the median age of all deaths was 81 years (IQR: 73-87 years). By district of residence, deceased cases were 85 (32.2%) from Nicosia, 81 (30.7%) from Limassol, 52 (19.7%) from Larnaka, 22 (8.3%) from Pafos, 16 (6.1%) from Ammochostos, and eight deaths (3%) occurred among cases reported either in the British bases or had a residence abroad, or information was not available.

The median time from date of sampling to death (all-causes) was 12 days (IQR: 6-20 days).

Figure 5b reports the number of all deaths among people with COVID-19 by date.




Figure 5a: Number of COVID-19 associated deaths in the Republic of Cyprus by date of death.




Figure 5b: Number of deaths (all causes) among COVID-19 cases in the Republic of Cyprus by date of death.





## Hospitalization and intensive care unit (ICU<sup>3</sup>) admissions<sup>4</sup>

In total, 5.1% (n = 1,603) of people with COVID-19 received hospital care, as of February 9<sup>th</sup>, 2021. The median age of hospitalized patients was 66 years (IQR: 52-78 years). Hospitalized cases were mainly males (n = 933; 58.3%).

The median age of patients still hospitalized as of 10<sup>th</sup> of February, 2021 (n = 98, including those discharged on that day) is 67 years (IQR: 53-75 years), 52% (n = 51) are males, and 44.9% (n = 44) are from Limassol district. Sixty-four cases (65.3%) still hospitalised have comorbidities.

Figure 6a shows the total number of first hospital admissions by date.




Figure 6a: Number and 7-day moving average (last moving average on Feb 9<sup>th</sup>, 2021 is shown on the graph) of laboratory-confirmed COVID-19 cases by date of first hospital admission.

*\*Date of hospital admission was replaced with date of sampling for inpatients hospitalised prior to the beginning of the epidemic.*

**Recent data should be interpreted with caution due to the possibility that cases recently hospitalised could have not been notified.**

<sup>3</sup> Intensive care unit (ICU) refers only to the ICU in Limassol General Hospital and to the ICU in Nicosia General Hospital.

<sup>4</sup> Data on hospitalisation and ICU are provisional and should be interpreted with caution because delay in data reporting is likely; for the construction of the curve, people are no longer in an ICU/hospital the day next to the date of their discharge, death or transfer.



Figure 6b reports the number of individuals in hospital every day (excluding those in ICU).




Figure 6b: Number and 7-day moving average (last moving average on Feb 10<sup>th</sup>, 2021 is shown on the graph) of laboratory-confirmed COVID-19 cases in hospital every day (excluding those in ICU).

*Patients in hospital for only part of the day (e.g. transferred, dead) are included.*

***Recent data should be interpreted with caution due to the possibility that cases recently hospitalised could have not been notified.***



Overall, 166 cases (10.4% of all hospitalized patients) have been admitted to ICU. The median age of patients ever admitted to ICU was 70 years (IQR: 62-76 years). ICU patients are mainly male (n = 110; 66.3%). The overall median length of stay in ICU (for all ICU cases) was 15 days (IQR: 8-25 days).

As of February 10<sup>th</sup>, 2021 and based on records which rapidly evolve, 23 cases are still in ICU (including deaths / discharged on that day). The median age of current ICU patients is 70 (IQR: 60-76) years and 13 (56.5%) are females. Twelve (52.2%) patients currently in ICU have pre-existing conditions.

The number of cases currently in ICU is 2.6 per 100,000 population (as of February 10<sup>th</sup>, 2021, including deaths/discharged on that day).

A total of 153 ICU patients (92.2% of all ICU patients) have been intubated - currently there are 23 patients intubated in ICU (including deaths/discharged/extubated on that day).

Figure 7 shows the number of patients in ICU, by day and intubation status.




Figure 7: Number of laboratory-confirmed COVID-19 cases in ICU by date and intubation.

*\*Date of discharge/transfer/death included*

**Recent data should be interpreted with caution due to the possibility that cases recently admitted in ICU or intubated could have not been notified/recorded.**



## Recovered/released

As of February 10<sup>th</sup>, 2021, among cases alive, 90.7% (n = 28,518) of COVID-19 cases have recovered/released from isolation<sup>5</sup>; of which 1,315 (4.6%) tested negative two consecutive times, and 27,203 (95.4%) have been released as per later guidelines<sup>6</sup>.

---

<sup>5</sup> For symptomatic cases, or for cases hospitalised, a COVID-19 case can be considered cured after the resolution of symptoms and two negative tests for SARS-CoV-2 at 24-hour interval at least.

For asymptomatic cases, or for persons isolated at home, the negative tests to document virus clearance should be obtained at a minimum of 14 days after the initial positive test (end of the quarantine period).

Novel coronavirus (SARS-CoV-2). Discharge criteria for confirmed COVID-19 cases- When is it safe to discharge COVID-19 cases from the hospital or end home isolation? - Technical Report, 10 March 2020. Stockholm: ECDC; 2020.

<sup>6</sup> May – November 2020: A person is released 21 days after the initial diagnosis, if he/she has a positive test 14 days after the initial diagnosis and remains in isolation for one more week without being further tested.

Since 8th November 2020: A person who remained asymptomatic throughout his/her follow-up is released 14 days after initial diagnosis (date of sampling) and General Practitioner (GP) confirmation; a person with mild/moderate symptoms is released in absence of fever AND improvement of other symptoms (confirmed by GP) AND 14 days after onset of symptoms; cases requiring hospitalisation (severe cases), immunosuppressed individuals, residents and staff of vulnerable groups (i.e., nursing homes, refugee camps, prisons) are released in absence of fever AND 20 days after onset of symptoms. The criterion of two consecutive negative SARS-CoV-2 PCR results from respiratory samples, taken at least 24 hours apart, is recommended for immunosuppressed and critically ill COVID-19 patients.



## **Acknowledgments**

We would like to thank the KIOS Center of Excellence of the University of Cyprus for the development of the ICT system to support data management.

## Appendix

Table A1: Characteristics of cases by origin of infection diagnosed in the last 14 days (27 Jan – 09 Feb, 2021).

Characteristics	Total (n = 1536)		Imported (n = 16)		Locally-acquired (n = 1520)	
	N	%	N	%	N	%
Sex						
Male	762	49.6	8	50.0	754	49.6
Female	773	50.3	8	50.0	765	50.3
Unknown	1	0.1	0	0.0	1	0.1
Age groups (years)						
0-9	111	7.2	1	6.3	110	7.2
10-19	142	9.2	2	12.5	140	9.2
20-29	258	16.8	3	18.8	255	16.8
30-39	295	19.2	5	31.3	290	16.8
40-49	251	16.3	1	6.3	250	19.1
50-59	214	13.9	4	25.0	210	16.4
60-69	154	10.0	0	0.0	154	13.8
70-79	81	5.3	0	0.0	81	10.1
80+	30	2.0	0	0.0	30	5.3
Unknown	1	0.1	0	0.0	1	2.0
Median age in years (IQR)	38 (25-54)		32 (23-48)		38 (25-54)	
District						
Ammochostos	51	3.3	4	25.0	47	3.1
Larnaka	181	11.8	7	43.8	174	11.4
Limassol	835	54.4	3	18.8	832	54.7
Nicosia	370	24.1	2	12.5	368	24.2
Pafos	61	4.0	0	0.0	61	4.0
Other	38	2.5	0	0.0	38	2.5
Symptoms at diagnosis						
Yes	992	64.6	8	50.0	984	64.7
No	544	35.4	8	50.0	536	35.3
Unknown	0	0.0	0	0.0	0	0.0



Figure A1. 14-day cumulative diagnosis rate (per 100,000 population) by age group for the last 14 days.






Figure A2: 14-day cumulative diagnosis rate (per 100,000 population) by district for the last 14 days.








Figure A3A: Number of cases by origin of infection and by date of sample collection in the last 14 days (27 Jan – 09 Feb, 2021).






Figure A3B: Number and 7-day moving average of locally-acquired cases, by date of sample collection and district in the last 14 days (27 Jan – 09 Feb, 2021).

