



# **Coronavirus Disease 2019 (COVID-19)**

## **National Surveillance Report as of 12/01/2021**

Date of publication: 15/01/2021

Produced by the Epidemiological Surveillance Unit of the Ministry of Health

Contributors: Annalisa Quattrocchi, Ioannis Mamais, Valentinos Silvestros, Anna Demetriou (Health Monitoring Unit), Maria Athanasiadou (Health Monitoring Unit), Theopisti Kyprianou (Health Monitoring Unit), Androulla Stylianou, Sotiroula Sotiriou, Fani Theofhanous, Christos Charalambous, Ioanna Gregoriou, Maria Koliou, Olga Kalakouta, Georgios Nikolopoulos

Scientific Committee: Constantinos Constantinou, Niki Paphitou, Georgios Nikolopoulos, Maria Koliou, George Panos, Eirini Christaki, Zoi – Dorothea Pana, Constantinos Tsioutis, Leondios Kostrikis, Peter Karayiannis, George Petrikkos, Petros Agathangelou, George Mixides

Suggested citation: Epidemiological Surveillance Unit of the Ministry of Health, Cyprus. National Situation Report. Coronavirus Disease 2019 (COVID-19), 15 January 2021.



## Summary

- As of January 12<sup>th</sup>, 2021, a total of 27,827 COVID-19 cases have been diagnosed, of which 165 died due to COVID-19 (case fatality risk: 0.6%).
- The 14-day cumulative diagnosis rate is 665.2 per 100,000 population.
- In the last 14 days (30 Dec, 2020 - 12 Jan, 2021), 5,907 cases were diagnosed.
  - The median age was 39 years (Interquartile range - IQR: 25-56 years); 53.2% were female (n = 3,140), 46.7% male (n = 2,756), and for 11 cases (0.2%) sex is unknown at the moment.
  - By place of exposure: 1% (n = 60) were imported and 99% (n = 5,847) were locally-acquired.
- As of January 13<sup>th</sup>, 2021, 209 people were still hospitalized (currently notified). The median age of patients still hospitalized is 71 years (IQR: 62-79 years), 58.9% (n = 123) are males, and 30.1% (n = 63) are from Nicosia district.
- Of 38 cases in intensive care units (ICU), who are currently notified, 28 are intubated. The median age of current ICU patients is 67 (IQR: 61-77) years and 24 (63.2%) are males. Twenty-five (65.8%) patients currently in ICU have pre-existing conditions.



## Epidemiological surveillance in the Republic of Cyprus

Analyses are based on laboratory-confirmed cases notified to the Epidemiological Surveillance Unit of the Ministry of Health.

As of January 12<sup>th</sup>, 2021, 27,827 cases of coronavirus disease 2019 (COVID-19) have been diagnosed (laboratory-confirmed) (Figure 1, 2 and 3).

In the last 14 days (30 Dec, 2020 - 12 Jan, 2021), 5,907 cases have been diagnosed. The 14-day cumulative diagnosis rate of COVID-19 (per 100,000 population)<sup>1</sup>, a measure which reflects the number of active COVID-19 cases in the population (prevalence)<sup>2</sup>, is 665.2 per 100,000 population (Figure 3).




Figure 1: Number and 7-day moving average (last moving average on Jan 12<sup>th</sup>, 2021, is shown on the graph) of laboratory-confirmed COVID-19 cases in Cyprus since 03/03/2020 by date of sample collection\*.

\* when date of sample collection was not available, laboratory result date or hospitalisation date was used.

<sup>1</sup>Population denominator for end of year 2019, available at [https://www.mof.gov.cy/mof/cystat/statistics.nsf/populationcondition\\_21main\\_en/populationcondition\\_21main\\_en?OpenForm&sub=1&sel=4](https://www.mof.gov.cy/mof/cystat/statistics.nsf/populationcondition_21main_en/populationcondition_21main_en?OpenForm&sub=1&sel=4)

<sup>2</sup>Coronavirus disease 2019 (COVID-19) pandemic: increased transmission in the EU/EEA and the UK – seventh update, 25 March 2020. Stockholm: ECDC; 2020. <https://www.ecdc.europa.eu/sites/default/files/documents/RRA-seventh-update-Outbreak-of-coronavirus-disease-COVID-19.pdf>




Figure 2: Number of laboratory-confirmed COVID-19 cases in Cyprus since 03/03/2020 by date of symptoms onset.

**Recent data should be interpreted with caution due to the possibility that cases with date of onset within the reporting period have not yet been diagnosed.**




Figure 3: COVID-19 14-day cumulative diagnosis rate per 100,000 population (proxy of COVID-19 prevalence).



## Characteristics of cases diagnosed in the last 14 days

### Age, sex, and place of residence

Among 5,907 cases diagnosed since 30<sup>th</sup> December, 2020, 53.2% were female (n = 3,140), 46.7% male (n = 2,756), and for 11 cases (0.2%) sex is unknown at the moment.

By age group, cases included 863 infants, children and adolescents aged 0-19 years-old (14.6%), 3,852 adults aged 20-59 years (65.2%), 1,186 persons aged 60 years and older (20.1%), and for 6 cases (0.1%) age is unknown at the moment. Figure 4 shows the number of cases by 10-year age band and sex. The median age of all cases diagnosed in the last 14 days is 39 years (IQR: 25-56 years). The median age of adult cases ( $\geq 18$  years) is 43 years (IQR: 30-58 years).

The 14-day cumulative diagnosis rate (per 100,000 population) by age group is shown in figure A1 in Appendix.




Figure 4: Laboratory-confirmed COVID-19 cases in Cyprus, diagnosed in the last 14 days, by age group and sex.

Among all cases diagnosed in the last 14 days, 2,180 (36.9%) were reported in Nicosia district, 1,674 (28.3%) in Limassol, 1,332 (22.6%) in Larnaka, 327 (5.5%) in Pafos, 295 (5%) in Ammochostos, and 99 (1.7%) were reported either in the British bases or had a residence abroad, or information was not available. Figure A2 in Appendix shows the 14-day cumulative diagnosis rate (per 100,000 population) by district.



### Epidemiological link

In the last 14 days (30 Dec, 2020 – 12 Jan, 2021), of 5,907 cases sampled and notified, 1% (n = 60) were imported and 99% (n = 5,847) were locally-acquired. Table A1 and Figure A3 (A&B) in Appendix show the characteristics of cases by origin of infection and the daily number of locally-acquired cases by district in the recent period, respectively.

### Clinical features

In the last 14 days (30 Dec, 2020 – 12 Jan, 2021), 28.9% (n = 1,709) reported no symptoms at diagnosis and 71.1% (n = 4,198) reported at least one symptom.

### Pre-existing conditions

In the last 14 days (30 Dec, 2020 – 12 Jan, 2021), 1,608 cases (27.2%) reported at least one comorbidity.



## Deaths

Among cases, 165 COVID-19 associated deaths were reported in Cyprus until January 14<sup>th</sup>, 2021 (Case Fatality Risk - CFR: 0.6%).

The COVID-19 associated mortality is 18.6 per 100,000 population.

Deaths occurred in 109 men (66.1%) and 56 (33.9%) women; the median age of all COVID-19 associated deaths was 82 years (IQR: 74-87 years). By district of residence, deceased cases were 57 (34.6%) from Nicosia, 52 (31.5%) from Limassol, 28 (17%) from Larnaka, 13 (7.9%) from Ammochostos, 12 (7.3%) from Pafos and three deaths (1.8%) occurred among cases reported either in the British bases or had a residence abroad, or information was not available.

The median time from date of sampling to death (due to COVID-19) was 11 days (IQR: 6-18 days).

Figure 5a reports the number of COVID-19 associated deaths by date.

Among cases, 202 deaths (all causes) were reported in Cyprus until January 14<sup>th</sup>, 2021 (CFR: 0.7%).

The mortality (all-causes) for people with COVID-19 is 22.7 per 100,000 population.

Deaths occurred in 132 men (65.4%) and 70 (34.6%) women; the median age of all deaths was 82 years (IQR: 73-87 years). By district of residence, deceased cases were 66 (32.7%) from Limassol, 64 (31.7%) from Nicosia, 37 (18.3%) from Larnaka, 15 each (7.4%) from Ammochostos and Pafos, and five deaths (2.5%) occurred among cases reported either in the British bases or had a residence abroad, or information was not available.

The median time from date of sampling to death (all-causes) was 10 days (IQR: 5-17 days).

Figure 5b reports the number of all deaths among people with COVID-19 by date.




Figure 5a: Number of COVID-19 associated deaths in Cyprus by date of death.




Figure 5b: Number of deaths (all causes) among COVID-19 cases in Cyprus by date of death.





## Hospitalization and intensive care unit (ICU<sup>3</sup>) admissions<sup>4</sup>

In total, 4.6% (n = 1,271) of people with COVID-19 received hospital care, as of January 13<sup>th</sup>, 2021. The median age of hospitalized patients was 66 years (IQR: 52-79 years). Hospitalized cases were mainly males (n = 734; 57.8%).

The median age of patients still hospitalized as of 14<sup>th</sup> of January, 2021 (n = 209) is 71 years (IQR: 62-79 years), 58.9% (n = 123) are males, and 30.1% (n = 63) are from Nicosia district. One-hundred-thirty-four cases (64.1%) still hospitalised have comorbidities.

Figure 6a shows the total number of first hospital admissions by date.




Figure 6a: Number and 7-day moving average (last moving average on Jan 14<sup>th</sup>, 2021 is shown on the graph) of laboratory-confirmed COVID-19 cases by date of first hospital admission.

*\*Date of hospital admission was replaced with date of sampling for inpatients hospitalised prior to the beginning of the epidemic.*

**Recent data should be interpreted with caution due to the possibility that cases recently hospitalised could have not been notified.**

<sup>3</sup> Intensive care unit (ICU) refers only to the ICU in Limassol General Hospital and to the ICU in Nicosia General Hospital.

<sup>4</sup> Data on hospitalisation and ICU are provisional and should be interpreted with caution because delay in data reporting is likely; for the construction of the curve, people are no longer in an ICU/hospital the day next to the date of their discharge, death or transfer.



Figure 6b reports the number of individuals in hospital every day (excluding those in ICU).




Figure 6b: Number and 7-day moving average (last moving average on Jan 14<sup>th</sup>, 2021 is shown on the graph) of laboratory-confirmed COVID-19 cases in hospital every day (excluding those in ICU).

*Patients in hospital for only part of the day (e.g. transferred, dead) are included.*

***Recent data should be interpreted with caution due to the possibility that cases recently hospitalised could have not been notified.***



Overall, 125 cases (9.8% of all hospitalized patients) have been admitted to ICU. The median age of patients ever admitted to ICU was 69 years (IQR: 62-76 years). ICU patients are mainly male (n = 87; 69.6%). The overall median length of stay in ICU (for all ICU cases) was 11 days (IQR: 7-22 days).

As of January 14<sup>th</sup>, 2021 and based on records which rapidly evolve, 38 cases are still in ICU (including deaths / discharged on that day). The median age of current ICU patients is 67 (IQR: 61-77) years and 24 (63.2%) are males. Twenty-five (65.8%) patients currently in ICU have pre-existing conditions.

The number of cases currently in ICU is 4.3 per 100,000 population (as of January 13<sup>th</sup>, 2021, including deaths/discharged on that day).

A total of 113 ICU patients (90.4% of all ICU patients) have been intubated - currently there are 28 patients intubated in ICU (including deaths/discharged on that day). Figure 7 shows the number of patients in ICU, by day and intubation status.




Figure 7: Number of laboratory-confirmed COVID-19 cases in ICU by date and intubation.

*\*Date of discharge/transfer/death included*

***Recent data should be interpreted with caution due to the possibility that cases recently admitted in ICU or intubated could have not been notified/recorded.***



## Recovered/released

As of January 13<sup>th</sup>, 2021, among cases alive, 67.8% (n = 18,743) of COVID-19 cases have recovered/released from isolation<sup>5</sup>; of which 1,314 (7%) tested negative two consecutive times, and 17,429 (93%) have been released as per later guidelines<sup>6</sup>.

*More updated information on recovery, based on new guidelines that do not require testing, will be available in next reports.*

---

<sup>5</sup> For symptomatic cases, or for cases hospitalised, a COVID-19 case can be considered cured after the resolution of symptoms and two negative tests for SARS-CoV-2 at 24-hour interval at least.

For asymptomatic cases, or for persons isolated at home, the negative tests to document virus clearance should be obtained at a minimum of 14 days after the initial positive test (end of the quarantine period).

Novel coronavirus (SARS-CoV-2). Discharge criteria for confirmed COVID-19 cases- When is it safe to discharge COVID-19 cases from the hospital or end home isolation? - Technical Report, 10 March 2020. Stockholm: ECDC; 2020.

<sup>6</sup> May – November 2020: A person is released 21 days after the initial diagnosis, if he/she has a positive test 14 days after the initial diagnosis and remains in isolation for one more week without being further tested.

Since 8th November 2020: A person who remained asymptomatic throughout his/her follow-up is released 14 days after initial diagnosis (date of sampling) and General Practitioner (GP) confirmation; a person with mild/moderate symptoms is released in absence of fever AND improvement of other symptoms (confirmed by GP) AND 14 days after onset of symptoms; cases requiring hospitalisation (severe cases), immunosuppressed individuals, residents and staff of vulnerable groups (i.e., nursing homes, refugee camps, prisons) are released in absence of fever AND 20 days after onset of symptoms. The criterion of two consecutive negative SARS-CoV-2 PCR results from respiratory samples, taken at least 24 hours apart, is recommended for immunosuppressed and critically ill COVID-19 patients.



## **Acknowledgments**

We would like to thank the KIOS Center of Excellence of the University of Cyprus for the development of the ICT system to support data management.

## Appendix

Table A1: Characteristics of cases by origin of infection diagnosed in the last 14 days (30 Dec, 2020 – 12 Jan, 2021).

Characteristics	Total (n = 5907)		Imported (n = 60)		Locally-acquired (n = 5847)	
	N	%	N	%	N	%
Sex						
Male	2756	46.7	29	48.3	2727	46.6
Female	3140	53.2	30	50.0	3110	53.2
Unknown	11	0.2	1	1.7	10	0.2
Age groups (years)						
0-9	315	5.3	0	0.0	315	5.4
10-19	548	9.3	4	6.7	544	9.3
20-29	1102	18.7	19	31.7	1083	18.5
30-39	1064	18.0	9	15.0	1055	18.5
40-49	790	13.4	7	11.7	783	13.4
50-59	896	15.2	13	21.7	883	15.1
60-69	624	10.6	6	10.0	618	10.6
70-79	352	6.0	1	1.7	351	6.0
80+	210	3.6	1	1.7	209	3.6
Unknown	6	0.1	0	0.0	6	0.1
Median age in years (IQR)	39 (25-56)		37 (24-54)		39 (25-56)	
District						
Ammochostos	295	5.0	13	21.7	282	4.8
Larnaka	1332	22.5	16	26.7	1316	22.5
Limassol	1674	28.3	10	16.7	1664	28.5
Nicosia	2180	36.9	8	13.3	2172	37.1
Pafos	327	5.5	10	16.7	317	5.4
Other	99	1.7	3	5.0	96	1.6
Symptoms at diagnosis						
Yes	4,198	71.1	25	41.7	4173	71.4
No	1709	28.9	35	58.3	1674	28.6
Unknown	0	0.0	0	0.0	0	0.0



Figure A1. 14-day cumulative diagnosis rate (per 100,000 population) by age group for the last 14 days.






Figure A2: 14-day cumulative diagnosis rate (per 100,000 population) by district for the last 14 days.








Figure A3A: Number of cases by origin of infection and by date of sample collection in the last 14 days (30 Dec, 2020 – 12 Jan, 2021).






Figure A3B: Number and 7-day moving average of locally-acquired cases, by date of sample collection and district in the last 14 days (30 Dec, 2020 – 12 Jan, 2021).

