



# **Coronavirus Disease 2019 (COVID-19)**

## **National Surveillance Report as of 17/01/2022**

Date of publication: 21/01/2022

Produced by the Epidemiological Surveillance Unit of the Ministry of Health

Contributors: Annalisa Quattrocchi, Ioannis Mamais, Valentinos Silvestros, Anna Demetriou (Health Monitoring Unit), Maria Athanasiadou (Health Monitoring Unit), Theopisti Kyprianou (Health Monitoring Unit), Androulla Stylianos, Sotiroula Sotiriou, Fani Theofhanous, Christos Charalambous, Ioanna Gregoriou, Maria Koliou, Olga Kalakouta, Georgios Nikolopoulos

Scientific Committee: Constantinos Constantinou, Niki Paphitou, Georgios Nikolopoulos, Maria Koliou, George Panos, Eirini Christaki, Zoi – Dorothea Pana, Constantinos Tsioutis, Linos Hadjihannas, Christos Petrou, Peter Karayiannis, George Petrikkos, Petros Agathangelou, Constantinos Phellas, George Mixides

Suggested citation: Epidemiological Surveillance Unit of the Ministry of Health, Cyprus. National Situation Report. Coronavirus Disease 2019 (COVID-19), 21 January 2022.



## Summary

- As of January 17<sup>th</sup>, 2022, 223,261 COVID-19 cases have been diagnosed, of which 697 died due to COVID-19 (case fatality risk: 0.3%).
- In the last 14 days (since 4<sup>th</sup> January, 2021), 46,487 cases were diagnosed. The 14-day cumulative diagnosis rate is 5235 per 100,000 population.
  - The median age was 32 years (IQR: 20-46 years); sex information was available for 38,388 (82.6%) cases, of which 51.7% were females (n = 19,856), and 48.3% males (n = 18,532).
  - By place of exposure, information was available for 46,047 (99.1%) cases, of which 1.6% (n = 749) were imported and 98.4% (n = 45,298) were locally-acquired.
- As of January 19<sup>th</sup>, 2022, 263 people were still hospitalized (currently notified and among those diagnosed till January 17<sup>th</sup> including those discharged/who died on January 19<sup>th</sup>). The median age of patients still hospitalized due to COVID-19 (n = 263, including those discharged on that day) is 66 years (IQR: 55-74 years), 59.2% (n = 132 out of 223 with available information) are males, and 33.3% (n = 65 out of 195 with information on district) are from Limassol district. Forty-four cases (20.3%) still hospitalised have comorbidities (out of 217).
- As January 19<sup>th</sup>, 2022, of 44 cases in intensive care units (ICU), who are currently notified and diagnosed till January 17<sup>th</sup> including those transferred/who died on January 19<sup>th</sup>, 42 (95.5%) are intubated. The median age of current ICU patients is 65 (IQR: 58-71) years and 24 (58.5%) are males (out of 41 cases with available information). Fifteen (35.7%) ICU patients have comorbidities (out of 42).
- Over the last 14 days, 132,832 RT PCR and 1,538,625 rapid antigen tests have been performed (14,958.6 RT PCR and 173,268.6 rapid antigen tests per 100,000 population).



## Epidemiological surveillance in the Republic of Cyprus

Analyses are based on laboratory-confirmed cases notified to the Epidemiological Surveillance Unit of the Ministry of Health.

As of January 17<sup>th</sup>, 2022, 223,261 cases of coronavirus disease 2019 (COVID-19) have been diagnosed (laboratory-confirmed) (Figure 1, 2 and 3).

In the last 14 days (4 – 17 January 2022), 46,487 cases have been diagnosed. The 14-day cumulative diagnosis rate of COVID-19 (per 100,000 population)<sup>1</sup>, a measure which reflects the number of active COVID-19 cases in the population (prevalence)<sup>2</sup>, is 5,235 per 100,000 population (Figure 3).




Figure 1: Number and 7-day moving average (last moving average on January 17<sup>th</sup>, 2022, is shown on the graph) of laboratory-confirmed COVID-19 cases in the Republic of Cyprus since 03/03/2020 by date of sample collection\*.

\* when date of sample collection was not available, laboratory result date or hospitalisation date was used.

<sup>1</sup>Population denominator for end of year 2019, available at [https://www.mof.gov.cy/mof/cystat/statistics.nsf/populationcondition\\_21main\\_en/populationcondition\\_21main\\_en?OpenForm&sub=1&sel=4](https://www.mof.gov.cy/mof/cystat/statistics.nsf/populationcondition_21main_en/populationcondition_21main_en?OpenForm&sub=1&sel=4)

<sup>2</sup>Coronavirus disease 2019 (COVID-19) pandemic: increased transmission in the EU/EEA and the UK – seventh update, 25 March 2020. Stockholm: ECDC; 2020. <https://www.ecdc.europa.eu/sites/default/files/documents/RRA-seventh-update-Outbreak-of-coronavirus-disease-COVID-19.pdf>




Figure 2: Number of laboratory-confirmed COVID-19 cases in the Republic of Cyprus since 03/03/2020 by date of symptoms onset.  
**Recent data should be interpreted with caution due to the possibility that cases with date of onset within the reporting period have not yet been diagnosed.**




Figure 3: COVID-19 14-day cumulative diagnosis rate per 100,000 population (proxy of COVID-19 prevalence).

## Characteristics of cases diagnosed in the last 14 days

### Age, sex, and place of residence

Among 46,487 cases diagnosed since January 4<sup>th</sup>, 2021, sex information was available for 38,388 (82.6%) cases, of which 51.7% were females (n = 19,856), and 48.3% males (n = 18,532).

By age group, cases included 11,297 individuals aged 0-19 years-old (24.3%), 30,481 individuals aged 20-59 years (65.6%), and 4,707 individuals aged 60 years and older (10.1%), while for two cases age was not known. Figure 4 shows the number of cases by 10-year age band and sex. The median age of all cases diagnosed in the last 14 days is 32 years (IQR: 20-46 years). The median age of adult cases ( $\geq 18$  years) is 37 years (IQR: 28-50 years).

The 14-day cumulative diagnosis rate (per 100,000 population) by age group is shown in figure A1 in the Appendix.




Figure 4: Laboratory-confirmed COVID-19 cases in the Republic of Cyprus, diagnosed in the last 14 days, by age group and sex (including only cases with sex information available).

Among all cases diagnosed in the last 14 days, district information was available for 37,819 (81.4%) cases, of which 15,121 (40%) were reported in Nicosia district, 10,783 (28.5%) in Limassol, 6,166 (16.3%) in Larnaca, 3,565 (9.4%) in Pafos, and 2,148 (5.7%) in Ammochostos and 36 (0.1) cases had residence abroad. Figure A2 in the Appendix shows the 14-day cumulative diagnosis rate (per 100,000 population) by district.



### Epidemiological link

In the last 14 days (4 – 17 January 2022), epidemiological link information was available for 46,047 (99.1%) cases, of which 1.6% (n = 749) were imported and 98.4% (n = 45,298) were locally-acquired.

### Clinical features

In the last 14 days (4 – 17 January 2022), information was available for 42,754 (92%) cases of which 38.3% (n = 16,376) reported no symptoms and 61.7% (n = 26,378) reported at least one symptom.

### Pre-existing conditions

In the last 14 days (4 – 17 January 2022), information was available for 45,111 cases (97%) of which 2,651 cases (5.9%) reported at least one comorbidity.



## Deaths

Among cases diagnosed until January 17<sup>th</sup>, 2022, 697 COVID-19 associated deaths were reported in the Republic of Cyprus until January 19<sup>th</sup>, 2022 (Case Fatality Risk - CFR: 0.3%).

The COVID-19 associated mortality is 78.5 per 100,000 population.

Deaths occurred in 439 men (63%) and 258 (37%) women; the median age of all COVID-19 associated deaths was 77 years (IQR: 69-85 years). By district of residence, deceased cases were 239 (34.3%) from Limassol, 214 (30.7%) from Nicosia, 130 (18.7%) from Larnaca, 64 (9.2%) from Pafos, 40 (5.7%) from Ammochostos, and 10 deaths (1.4%) occurred among cases for which district information was not available or who had a residence abroad.

The median time from date of sampling to death (due to COVID-19) was 14 days (IQR: 8-23 days).

Figure 5a reports the number of COVID-19 associated deaths by date.

Among cases diagnosed until January 17<sup>th</sup>, 2022, 793 COVID-19 associated deaths were reported in the Republic of Cyprus until January 19<sup>th</sup>, 2022 (CFR: 0.4%).

The mortality (all-causes) for people with COVID-19 is 89.3 per 100,000 population.

Deaths occurred in 501 men (63.2%) and 292 (36.8%) women; the median age of all deaths was 77 years (IQR: 69-85 years). By district of residence, deceased cases were 279 (35.2%) from Limassol, 244 (30.8%) from Nicosia, 145 (18.3%) from Larnaca, 72 (9.1%) from Pafos, 43 (5.4%) from Ammochostos, and 10 deaths (1.2%) occurred among cases for which district information was not available or were reported among cases who had a residence abroad.

The median time from date of sampling to death (all-causes) was 14 days (IQR: 7-23 days).

Figure 5b reports the number of all deaths among people with COVID-19 by date.




Figure 5a: Number of COVID-19 associated deaths in the Republic of Cyprus by date of death.




Figure 5b: Number of deaths (all causes) among COVID-19 cases in the Republic of Cyprus by date of death.



## Hospitalization and intensive care unit (ICU<sup>3</sup>) admissions<sup>4</sup>

In total, 3.1% (n = 7,010) of people with COVID-19 received hospital care, as of January 19<sup>th</sup>, 2022. The median age of hospitalized patients was 60 years (IQR: 47-73 years). Excluding 70 cases (1%) for which sex information is not available, hospitalized cases were mainly males (n = 3,899; 56.2%).

The median age of patients diagnosed till January 17<sup>th</sup> and still hospitalized, due to COVID-19, as of January 19<sup>th</sup>, 2022 (n = 263, including those discharged/who died on that day) is 66 years (IQR: 55-74 years), 59.2% (n = 132 out of 223 with available information) are males, and 33.3% (n = 65 out of 195 with information on district) are from Limassol district. Forty-four cases (20.3%) still hospitalised have comorbidities (out of 217 with available information).

Figure 6a shows the total number of first hospital admissions by date.




Figure 6a: Number and 7-day moving average (last moving average on January 19<sup>th</sup>, 2022 is shown on the graph) of laboratory-confirmed COVID-19 cases by date of first hospital admission, since 03/03/2020.

*\*Date of hospital admission was replaced with date of sampling for inpatients hospitalised prior to the beginning of the epidemic.*

**Recent data should be interpreted with caution due to the possibility that cases recently hospitalised could have not been notified.**

<sup>3</sup> Intensive care unit (ICU) refers only to the ICU in Limassol General Hospital and to the ICU in Nicosia General Hospital.

<sup>4</sup> Data on hospitalisation and ICU are provisional and should be interpreted with caution because delay in data reporting is likely; for the construction of the curve, people are no longer in an ICU/hospital the day next to the date of their discharge, death or transfer.



Figure 6b reports the number of individuals in hospital every day (excluding those in ICU).




Figure 6b: Number and 7-day moving average (last moving average on January 19<sup>th</sup>, 2022 is shown on the graph) of laboratory-confirmed COVID-19 cases in hospital every day, since 03/03/2020, excluding those in ICU.

*Patients in hospital for only part of the day (e.g. transferred, dead) are included.*

**Recent data should be interpreted with caution due to the possibility that cases recently hospitalised could have not been notified.**



Overall, 771 cases (11% of all hospitalized patients) have been admitted to ICU. The median age of patients ever admitted to ICU was 66 years (IQR: 58-73 years). ICU patients were mainly males (n = 485, 62.9%). The overall median length of stay in ICU (for all ICU cases) was 12 days (IQR: 7-20 days).

As of January 19<sup>th</sup>, 2022, and based on records which rapidly evolve, of those diagnosed till January 17<sup>th</sup>, 44 cases are still in ICU (including deaths / transfers on January 19<sup>th</sup>). The median age of current ICU patients is 65 (IQR: 58-71) years and 24 (58.5%) are males (out of 41 cases with available information). Fifteen (35.7%) ICU patients have comorbidities (out of 42 with available information).

The number of cases currently in ICU is 5 per 100,000 population (as of January 19<sup>th</sup>, 2022, including deaths/discharged on that day).

A total of 735 ICU patients (95.3% of all ICU patients) have been intubated - currently 42 (95.5%) patients in ICU are intubated (including deaths/discharged/extubated on that day).

Figure 7 shows the number of patients in ICU, by day and intubation status.




Figure 7: Number of laboratory-confirmed COVID-19 cases in ICU by date and intubation.

*\*Date of discharge/transfer/death included*

**Recent data should be interpreted with caution due to the possibility that cases recently admitted in ICU or intubated could have not been notified/recorded.**



## Recovered/released

As of January 17<sup>th</sup>, 2022, among cases alive, 90.2% (n = 200,534) of COVID-19 cases have recovered/released from isolation<sup>5,6,7,8</sup>.

---

<sup>5</sup> For symptomatic cases, or for cases hospitalised, a COVID-19 case can be considered cured after the resolution of symptoms and two negative tests for SARS-CoV-2 at 24-hour interval at least. For asymptomatic cases, or for persons isolated at home, the negative tests to document virus clearance should be obtained at a minimum of 14 days after the initial positive test (end of the quarantine period).

Novel coronavirus (SARS-CoV-2). Discharge criteria for confirmed COVID-19 cases- When is it safe to discharge COVID-19 cases from the hospital or end home isolation? - Technical Report, 10 March 2020. Stockholm: ECDC; 2020.

<sup>6</sup> May – November 2020: A person is released 21 days after the initial diagnosis, if he/she has a positive test 14 days after the initial diagnosis and remains in isolation for one more week without being further tested.

Since 8<sup>th</sup> November 2020: A person who remained asymptomatic throughout his/her follow-up is released 14 days after initial diagnosis (date of sampling) and General Practitioner (GP) confirmation; a person with mild/moderate symptoms is released in absence of fever AND improvement of other symptoms (confirmed by GP) AND 14 days after onset of symptoms; cases requiring hospitalisation (severe cases), immunosuppressed individuals, residents and staff of vulnerable groups (i.e., nursing homes, refugee camps, prisons) are released in absence of fever AND 20 days after onset of symptoms. The criterion of two consecutive negative SARS-CoV-2 PCR results from respiratory samples, taken at least 24 hours apart, is recommended for immunosuppressed and critically ill COVID-19 patients.

<sup>7</sup> Since 4<sup>th</sup> August 2021: Asymptomatic individuals are discharged from self-isolation 10 days after testing positive for SARS-CoV-2. Individuals with mild/moderate symptoms are discharged from self-isolation 10 days after the onset of symptoms, provided that the patient has no fever for at least 3 days (without use of antipyretics) and has improvement of all other symptoms.

<sup>8</sup> Since 17<sup>th</sup> January 2022: SARS-CoV-2 positive cases are discharged 10 days after testing positive for the virus. In cases where the GP considers that the isolation of the case should continue due to persistent symptoms (fever and no improvement of the remaining symptoms two days before release) or in the case of immunosuppressed patients, the period of isolation may be extended.



## **Acknowledgments**

We would like to thank the KIOS Center of Excellence of the University of Cyprus for the development of the ICT system to support data management.

## Appendix

Figure A1. 14-day cumulative diagnosis rate (per 100,000 population) by age group.






Figure A2: 14-day cumulative diagnosis rate (per 100,000 population) by district for the last 14 days.

