



# **Coronavirus Disease 2019 (COVID-19)**

## **National Surveillance Report as of 26/07/2021**

Date of publication: 30/07/2021

Produced by the Epidemiological Surveillance Unit of the Ministry of Health

Contributors: Annalisa Quattrocchi, Ioannis Mamais, Valentinos Silvestros, Anna Demetriou (Health Monitoring Unit), Maria Athanasiadou (Health Monitoring Unit), Theopisti Kyprianou (Health Monitoring Unit), Androulla Stylianou, Sotiroula Sotiriou, Fani Theofhanous, Christos Charalambous, Ioanna Gregoriou, Maria Koliou, Olga Kalakouta, Georgios Nikolopoulos

Scientific Committee: Constantinos Constantinou, Niki Paphitou, Georgios Nikolopoulos, Maria Koliou, George Panos, Eirini Christaki, Zoi – Dorothea Pana, Constantinos Tsioutis, Peter Karayiannis, George Petrikkos, Petros Agathangelou, Constantinos Phellas, George Mixides

Suggested citation: Epidemiological Surveillance Unit of the Ministry of Health, Cyprus. National Situation Report. Coronavirus Disease 2019 (COVID-19), 30 July 2021.



## Summary

- As of July 26<sup>th</sup>, 2021, a total of 98,144 COVID-19 cases have been diagnosed, of which 415 died due to COVID-19 (case fatality risk: 0.4%).
- In the last 14 days (13 – 26 July, 2021), 12,463 cases were diagnosed. The 14-day cumulative diagnosis rate is 1403.5 per 100,000 population.
  - The median age was 28 years (IQR: 19-41 years); sex information was available for 8,428 (67.6%) cases, of which, 52.2% were males (n = 4,404), and 47.8% females (n = 4,024).
  - By place of exposure, information was available for 624 (5%) cases, of which 6.7% (n = 42) were imported and 93.3% (n = 582) were locally-acquired.
- As of July 28<sup>th</sup>, 2021, 255 people were still hospitalized (currently notified and among those diagnosed till July 26<sup>th</sup>). The median age of patients still hospitalized due to COVID-19 (n = 255, including those discharged on that day) is 57 years (IQR: 45-69 years), 55.4% (n = 126 out of 237 with available sex information) are males, and 31.9% (n = 51 out of 160 with information on district) are from Limassol district. Nineteen cases (25.3%) still hospitalised have comorbidities (out of 75 with available information).
- Of 35 cases in intensive care units (ICU), who are currently notified and diagnosed till July 26<sup>th</sup>, all are intubated. The median age of current ICU patients is 61 (IQR: 52-70) years and 20 (57.1%) are males.
- Over the last 14 days, 131,243 RT PCR and 883,986 rapid antigen tests have been performed (14,779.6 RT PCR and 99,548 rapid antigen tests per 100,000 population).

## Epidemiological surveillance in the Republic of Cyprus

Analyses are based on laboratory-confirmed cases notified to the Epidemiological Surveillance Unit of the Ministry of Health.

As of July 26<sup>th</sup>, 2021, 98,144 cases of coronavirus disease 2019 (COVID-19) have been diagnosed (laboratory-confirmed) (Figure 1, 2 and 3).

In the last 14 days (13 – 26 July, 2021), 12,463 cases have been diagnosed. The 14-day cumulative diagnosis rate of COVID-19 (per 100,000 population)<sup>1</sup>, a measure which reflects the number of active COVID-19 cases in the population (prevalence)<sup>2</sup>, is 1403.5 per 100,000 population (Figure 3).




Figure 1: Number and 7-day moving average (last moving average on July 26<sup>th</sup>, 2021, is shown on the graph) of laboratory-confirmed COVID-19 cases in the Republic of Cyprus since 03/03/2020 by date of sample collection\*.

\* when date of sample collection was not available, laboratory result date or hospitalisation date was used.

<sup>1</sup>Population denominator for end of year 2019, available at [https://www.mof.gov.cy/mof/cystat/statistics.nsf/populationcondition\\_21main\\_en/populationcondition\\_21main\\_en?OpenForm&sub=1&sel=4](https://www.mof.gov.cy/mof/cystat/statistics.nsf/populationcondition_21main_en/populationcondition_21main_en?OpenForm&sub=1&sel=4)

<sup>2</sup>Coronavirus disease 2019 (COVID-19) pandemic: increased transmission in the EU/EEA and the UK – seventh update, 25 March 2020. Stockholm: ECDC; 2020. <https://www.ecdc.europa.eu/sites/default/files/documents/RRA-seventh-update-Outbreak-of-coronavirus-disease-COVID-19.pdf>




Figure 2: Number of laboratory-confirmed COVID-19 cases in the Republic of Cyprus since 03/03/2020 by date of symptoms onset.

**Recent data should be interpreted with caution due to the possibility that cases with date of onset within the reporting period have not yet been diagnosed.**




Figure 3: COVID-19 14-day cumulative diagnosis rate per 100,000 population (proxy of COVID-19 prevalence).

## Characteristics of cases diagnosed in the last 14 days

### Age, sex, and place of residence

Among 12,463 cases diagnosed since July 13<sup>th</sup>, 2021, sex information was available for 8,428 (67.6%) cases, of which, 52.2% were males (n = 4,404), and 47.8% females (n = 4,024).

By age group, cases included 3,332 infants, children and adolescents aged 0-19 years-old (26.7%), 8,205 adults aged 20-59 years (65.8%), 922 persons aged 60 years and older (7.4%) and for four cases information is not currently available. Figure 4 shows the number of cases by 10-year age band and sex. The median age of all cases diagnosed in the last 14 days is 28 years (IQR: 19-41 years). The median age of adult cases ( $\geq 18$  years) is 33 years (IQR: 25-45 years).

The 14-day cumulative diagnosis rate (per 100,000 population) by age group is shown in figure A1 in the Appendix.




Figure 4: Laboratory-confirmed COVID-19 cases in the Republic of Cyprus, diagnosed in the last 14 days, by age group and sex (including only cases with sex information available).

Among all cases diagnosed in the last 14 days, district information was available for 9,120 (73.2%) cases, of which 2,919 (32%) were reported in Nicosia district, 2,833 (31.1%) in Limassol, 1,602 (17.6%) in Larnaka, 958 (10.5%) in Pafos, 792 (8.7%) in Ammochostos, and 16 (0.2%) were reported either in the British bases or had a residence abroad. Figure A2 in the Appendix shows the 14-day cumulative diagnosis rate (per 100,000 population) by district.



### Epidemiological link

In the last 14 days (13 – 26 July, 2021), of 12,463 cases sampled and notified, epidemiological link information was available for 624 (5%) cases, of which 6.7% (n = 42) were imported and 93.3% (n = 582) were locally-acquired.

### Clinical features

In the last 14 days (13– 26 July, 2021), information was available for 6,807 (54.6%) of cases of which 4.6% (n = 310) reported no symptoms and 95.4% (n = 6,497) reported at least one symptom.

### Pre-existing conditions

In the last 14 days (13– 26 July, 2021), information was available for 90 cases (0.7%) of which 21 cases (23.3%) reported at least one comorbidity.



## Deaths

Among cases diagnosed until 26<sup>th</sup> July 2021, 415 COVID-19 associated deaths were reported in the Republic of Cyprus until July 28<sup>th</sup>, 2021 (Case Fatality Risk - CFR: 0.4%).

The COVID-19 associated mortality is 46.7 per 100,000 population.

Deaths occurred in 271 men (65.3%) and 144 (34.7%) women; the median age of all COVID-19 associated deaths was 79 years (IQR: 70-86 years). By district of residence, deceased cases were 153 (36.9%) from Limassol, 142 (34.2%) from Nicosia, 71 (17.1%) from Larnaka, 27 (6.5%) from Pafos, 17 (4.1%) from Ammochostos, four deaths (1%) occurred among cases for which district information was not available and one (0.2%) death was reported among cases who had a residence abroad.

The median time from date of sampling to death (due to COVID-19) was 13 days (IQR: 7-22 days).

Figure 5a reports the number of COVID-19 associated deaths by date.

Among cases diagnosed until 26<sup>th</sup> July 2021, 493 deaths (all causes) were reported in the Republic of Cyprus until July 28<sup>th</sup>, 2021 (CFR: 0.5%).

The mortality (all-causes) for people with COVID-19 is 55.5 per 100,000 population.

Deaths occurred in 326 men (66.1%) and 167 (33.9%) women; the median age of all deaths was 79 years (IQR: 70-86 years). By district of residence, deceased cases were 186 (37.7%) from Limassol, 164 (33.3%) from Nicosia, 84 (17%) from Larnaka, 34 (6.9%) from Pafos, 20 (4.1%) from Ammochostos, four deaths (0.8%) occurred among cases for which district information was not available, and one (0.2%) death was reported among cases who had a residence abroad,.

The median time from date of sampling to death (all-causes) was 13 days (IQR: 7-22 days).

Figure 5b reports the number of all deaths among people with COVID-19 by date.




Figure 5a: Number of COVID-19 associated deaths in the Republic of Cyprus by date of death.




Figure 5b: Number of deaths (all causes) among COVID-19 cases in the Republic of Cyprus by date of death.





## Hospitalization and intensive care unit (ICU<sup>3</sup>) admissions<sup>4</sup>

In total, 4.6% (n = 4,522) of people with COVID-19 received hospital care, as of July 28<sup>th</sup>, 2021. The median age of hospitalized patients was 61 years (IQR: 48-73 years). Excluding 20 cases (0.4%) for which sex information is not available, hospitalized cases were mainly males (n = 2,590; 57.3%).

The median age of patients diagnosed till July 26<sup>th</sup> and still hospitalized, due to COVID-19, as of July 28<sup>th</sup>, 2021 (n = 255, including those discharged on that day) is 57 years (IQR: 45-69 years), 53.2% (n = 126 out of 237 with available sex information) are males, and 31.9% (n = 51 out of 160 with information on district) are from Limassol district. Nineteen cases (25.3%) still hospitalised have comorbidities (out of 75 with available information).

Figure 6a shows the total number of first hospital admissions by date.




Figure 6a: Number and 7-day moving average (last moving average on July 28<sup>th</sup>, 2021 is shown on the graph) of laboratory-confirmed COVID-19 cases by date of first hospital admission.

*\*Date of hospital admission was replaced with date of sampling for inpatients hospitalised prior to the beginning of the epidemic.*

***Recent data should be interpreted with caution due to the possibility that cases recently hospitalised could have not been notified.***

<sup>3</sup> Intensive care unit (ICU) refers only to the ICU in Limassol General Hospital and to the ICU in Nicosia General Hospital.

<sup>4</sup> Data on hospitalisation and ICU are provisional and should be interpreted with caution because delay in data reporting is likely; for the construction of the curve, people are no longer in an ICU/hospital the day next to the date of their discharge, death or transfer.



Figure 6b reports the number of individuals in hospital every day (excluding those in ICU).




Figure 6b: Number and 7-day moving average (last moving average on July 28<sup>th</sup>, 2021 is shown on the graph) of laboratory-confirmed COVID-19 cases in hospital every day (excluding those in ICU).

*Patients in hospital for only part of the day (e.g. transferred, dead) are included.*

***Recent data should be interpreted with caution due to the possibility that cases recently hospitalised could have not been notified.***



Overall, 453 cases (10% of all hospitalized patients) have been admitted to ICU. The median age of patients ever admitted to ICU was 67 years (IQR: 59-73 years). ICU patients were mainly male (n = 296; 65.3%). The overall median length of stay in ICU (for all ICU cases) was 12 days (IQR: 7-22 days).

As of July 28<sup>th</sup>, 2021 and based on records which rapidly evolve, of those diagnosed till July 26<sup>th</sup>, 35 cases are still in ICU (including deaths / discharged on that day). The median age of current ICU patients is 61 (IQR: 52-70) years and 20 (57.1%) are males.

The number of cases currently in ICU is 3.9 per 100,000 population (as of July 28<sup>th</sup>, 2021, including deaths/discharged on that day).

A total of 434 ICU patients (95.8% of all ICU patients) have been intubated - currently all patients in ICU are intubated (including deaths/discharged/extubated on that day). Figure 7 shows the number of patients in ICU, by day and intubation status.




Figure 7: Number of laboratory-confirmed COVID-19 cases in ICU by date and intubation.

*\*Date of discharge/transfer/death included*

***Recent data should be interpreted with caution due to the possibility that cases recently admitted in ICU or intubated could have not been notified/recorded.***



## Recovered/released

As of July 26<sup>th</sup>, 2021, among cases alive, 88.2% (n = 86,065) of COVID-19 cases have recovered/released from isolation<sup>5,6</sup>.

---

<sup>5</sup> For symptomatic cases, or for cases hospitalised, a COVID-19 case can be considered cured after the resolution of symptoms and two negative tests for SARS-CoV-2 at 24-hour interval at least.

For asymptomatic cases, or for persons isolated at home, the negative tests to document virus clearance should be obtained at a minimum of 14 days after the initial positive test (end of the quarantine period).

Novel coronavirus (SARS-CoV-2). Discharge criteria for confirmed COVID-19 cases- When is it safe to discharge COVID-19 cases from the hospital or end home isolation? - Technical Report, 10 March 2020. Stockholm: ECDC; 2020.

<sup>6</sup> May – November 2020: A person is released 21 days after the initial diagnosis, if he/she has a positive test 14 days after the initial diagnosis and remains in isolation for one more week without being further tested.

Since 8th November 2020: A person who remained asymptomatic throughout his/her follow-up is released 14 days after initial diagnosis (date of sampling) and General Practitioner (GP) confirmation; a person with mild/moderate symptoms is released in absence of fever AND improvement of other symptoms (confirmed by GP) AND 14 days after onset of symptoms; cases requiring hospitalisation (severe cases), immunosuppressed individuals, residents and staff of vulnerable groups (i.e., nursing homes, refugee camps, prisons) are released in absence of fever AND 20 days after onset of symptoms. The criterion of two consecutive negative SARS-CoV-2 PCR results from respiratory samples, taken at least 24 hours apart, is recommended for immunosuppressed and critically ill COVID-19 patients.



## **Acknowledgments**

We would like to thank the KIOS Center of Excellence of the University of Cyprus for the development of the ICT system to support data management.



## Appendix

Figure A1. 14-day cumulative diagnosis rate (per 100,000 population) by age group.




Figure A2: 14-day cumulative diagnosis rate (per 100,000 population) by district for the last 14 days.

